JUL 07, 2019

05 Agarose Gel Electrophoresis

Tjusls China¹

¹Tianjin University

Tjusls China
Tianjin University

GUIDELINES

[M] 0 Mass Percent

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.48rgzv6

Protocol Citation: Tjusls China 2019. 05 Agarose Gel Electrophoresis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.48rgzv6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jul 07, 2019

Last Modified: Jul 07, 2019

PROTOCOL integer ID:
25585

MATERIALS

MATERIALS

- ⊗ TAE (Tris-Acetate-EDTA) buffer, 1x **Contributed by users**
- ⊗ DNA samples **Contributed by users Catalog #/**
- ⊗ 1% Agarose gel **Contributed by users Catalog #/**
- ⊗ 10×green loading buffer **Contributed by users Catalog #/**

- 1 Use 1×TAE buffer to prepare 1% Agarose mix in a flask, then put it in the microwave and heat it as long as it takes to completely dissolve the Agarose.

- 2 Take out the conical flask, cool it in the wash basin to about 50°C. Add EB quickly, and then mix well. Pour the Agarose gel into gel tray and insert comb into slots. Let the gel solidify for 15-20min. Meanwhile, dilute the 10x green buffer to 1x and add to the DNA samples.

50 °C

00:15:00 ~ 00:20:00

- 3 Place the gel onto the electrophoresis apparatus ensuring that it is totally submerged in 1xTAE buffer. Carefully load each sample into its designated lane and 2ul DNA marker into a separate lane.

2 µL

- 4 Run at 120V for 20-25 min. If the sample have not completely separated, the time may be extended appropriately.

00:15:00 ~ 00:20:00

- 5 Check the gel using a gel imager or under UV light, then take a photo on Norma.