JUL 12, 2019

10 Affinity Chromatography

Tjusls China¹

¹Tianjin University

Chemistry Method Development Community

Tjusls China
Tianjin University

MATERIALS

MATERIALS

⊗ EDTA Contributed by users

⊗ 100ml GST Elution Buffer (5mM glutathione, 50mM Tris.HCl (pH8.0)) G-Biosciences Catalog #786-541

⊗ 5 columns Ni-NTA Resin, 1ml Spin Column G-Biosciences Catalog #786-944

⊗ GSH-Glo(TM) Glutathione Assay, 50ml Promega Catalog #V6912

⊗ Guanidine hydrochloride Bio Basic Inc. Catalog #GB0242.SIZE.100g

⊗ double distilled water (ddH2O) Contributed by users

⊗ Ni-binding buffer Contributed by users Catalog #/

⊗ Ni-Washing buffer Contributed by users Catalog #/

⊗ Ni-Elution buffer Contributed by users Catalog #/

⊗ Protein solution that needs to be purified Contributed by users Catalog #/

⊗ NiSO4 Contributed by users Catalog #/

⊗ GST columns Contributed by users Catalog #/

⊗ GST-washing buffer Contributed by users Catalog #/

⊗ GST-binding buffer Contributed by users Catalog #/

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.5cgg2tw

Protocol Citation: Tjusls China 2019. 10 Affinity Chromatography.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.5cgg2tw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jul 10, 2019

Last Modified: Jul 12, 2019

PROTOCOL integer ID:
25704

1 Lysis of the bacteria.

1. Resuspend the frozen cell paste as best you can in the Lysis Buffer using a 10 mL pipet or

whatever means necessary. Let this suspension incubate for 20 minutes at room temperature, or until the suspension becomes turbid and viscous due to release of the bacteria's genomic DNA.

10 mL 00:20:00

2. Smash the bacteria.

3. Centrifuge at 18,000 rpm in a big rotor for 40 minutes at 4°C. Save the pellet and the supernatant.

18000 rpm 00:40:00 4 °C

2 IMAC(Immobilized metal ion [affinity chromatography](#))

Procedure

1. Remove the Ni column from the 4°C refrigerator, which contains 20% alcohol. Wash the column with water for 10 minutes. Change to Ni-binding buffer for another 10 minutes and balance the Ni column.

4 °C

2. Add the protein solution to the column, let it flow naturally and bind to the column. Repeat until the medium turns gray (usually twice).

3. Add Ni-Washing buffer several times and let it flow. Take 5ul of wash solution and test with [Coomassie Brilliant Blue](#). Stop washing when it doesn't turn blue.

5 µL

4. Add Ni-Elution buffer several times. Check as above.

5. Collect the eluted proteins for further operation. Recycle columns: Wash with 0.2mM EDTA , let stand for 10 min then drain. Wash with 6M Guanidine hydrochloride, let stand for 10 min then drain. Wash with ddH₂O for three times. Fill up with NiSO₄, place on a shaker overnight at 4°C.

4 °C

GST (Affinity chromatography of glutathione transferas)

Procedure

1. Remove the GST column from the 4°C refrigerator. Wash the column with GST-binding buffer for 10 minutes to balance the GST column.

4 °C 00:10:00

2. Add the protein solution to the column, let it flow naturally and bind to the column.

3. Add GST-Washing buffer several times and let it flow. Take 5µl of wash solution and test with [Coomassie Brilliant Blue](#). Stop washing when it doesn't turn blue.

5 µL

4. Add GST-Elution buffer several times. Check as above.

5. Collect the eluted proteins for further operation. Recycle columns: Wash with GST-Elution buffer until [Coomassie Brilliant Blue](#) doesn't turn blue when tested. Wash with 6M Guanidine hydrochloride , let stand for 10 min then drain. Wash with ddH₂O for three times.

00:10:00