JUL 12, 2019

11 Concentration and fluid exchanging

Tjusls China¹

¹Tianjin University

Tjusls China
Tianjin University

GUIDELINES

1. Protein is an active molecule, ensuring it is always on the ice and clinging to the principle of freezing and melting quickly.
2. When the protein is concentrated, it should be inverted several times after each protein added and mixed.
3. When pipetting the protein with a pipette, tilt the concentrating tube at the angle of 45 degrees. The tip of the pipette tip is drawn against the corner of the upper chamber of the concentrating tube. Try not to touch the filter.

MATERIALS

MATERIALS

⊗ DTT P212121 Catalog #SV-DTT

⊗ EDTA Contributed by users

⊗ NaOH Contributed by users

⊗ 20 % ethanol Contributed by users Catalog #/

⊗ Concentrating tube Contributed by users Catalog #/

BEFORE START INSTRUCTIONS

1. The concentrating tubes has different pore sizes, and it is preferable to select a pore size smaller than one third of the molecular weight of the interest protein.
2. Pre-cooled to 4 °C before use the thermo centrifuge.

🌡 4 °C

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5e6g3he

Protocol Citation: Tjusls China 2019. 11 Concentration and fluid exchanging.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.5e6g3he>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jul 12, 2019

Last Modified: Jul 12, 2019

PROTOCOL integer ID:
25790

Concentrate

1. [The default concentrator is already inactive.](#)
2. Pre-cool the Thermo centrifuge to 4 °C in advance.

4 °C

3. Pour the protein solution to be concentrated into a concentrating tube, level it with the trim tube, place it in a Thermo centrifuge at 3400 rpm, stop at half an hour, pour off the effluent, and add the protein solution to the upper layer.

3400 rpm

4. If the protein is used for long crystals, it can be concentrated to less than 500 µl. According to the amount of protein and protein performance, it is determined that the protein is colloid rather than turbid liquid. If the protein is used for 24 ml molecular sieves, concentrate to within 1 ml.

5. Pipet the concentrated protein into a Ep tube with a 200 µl pipette (with a yellow tip).

200 µL

Disposal of the reuse of the concentrating tube

- 2
 1. Rinse the upper chamber of the used concentrator tube with distilled water several times, and add some hypersaline solution overnight at room temperature.
 2. Pour off the sodium hydroxide from the upper chamber, rinse several times with distilled water, and centrifuge 3000 rpm for 5 minutes with distilled water.

3000 rpm 00:05:00

3. Step 2 is repeated 3 times.

4. Add 20% ethanol or ddH₂O to the upper chamber and store at 4 °C.

4 °C

change the liquid

- 3 The steps are the same as the concentration, and the buffer is added without any what.