VERSION 1

JUL 30, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.qckdsuw

Protocol Citation: Roey Angel, Eva Petrova 2020. 16S Bacteria 338F-516P-805R BSA. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.qckdsuw>

MANUSCRIPT CITATION:
 Yu, Y., Lee, C., Kim, J., and Hwang, S. (2005). Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnol Bioeng* 89, 670–679. doi:10.1002/bit.20347.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 24, 2018

🌐 16S Bacteria 338F-516P-805R BSA V.1

Roey Angel¹, Eva Petrova¹

¹Soil and Water Research Infrastructure

SoWa RI Anaerobic and Molecular Microbiology (public)
 Tech. support email: eva.petrova@bc.cas.cz

Roey Angel

Soil and Water Research Infrastructure

ABSTRACT

Universal 16S rRNA probe-based-qPCR assay for bacteria.
 The primers and probe are taken from [Yu et al. \(2005\)](#).

CITATION

Yu Y, Lee C, Kim J, Hwang S (2005). Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnology and bioengineering*.

LINK
<http://dx.doi.org/10.1002/bit.20347>

ATTACHMENTS

[Introduction_QPCR_Strata gene.pdf](#) [AB_rt-QPCRguide.pdf](#)

MATERIALS

MATERIALS

iQ™ SYBR® Green Supermix BioRad Sciences Catalog #1708880

PROTOCOL integer ID:
12396

Keywords: qPCR, dual-labelled probe, 16S rRNA gene, bacteria

Primers and probe

1

Name	Type	Sequence	Target region
BAC338F	Forward	ACT CCT ACG GGA GGC AG	338-354
BAC516P ²	Probe	TGC CAG CAG CCG CGG TAA TA	516-536
BAC805R	Reverse	GAC TAC CAG GGT ATC TAA TC	785-805

1. Relative to *E. coli* 16S rRNA gene.
2. The probe must be dual-labelled either with 5'-6-FAM, 3'-BHQ1 or any other valid combination.

qPCR mixture

2

Reagent	Final concentration	1 tube (20 µl)	plate (20 µl x 100)
PCR H ₂ O		4.6	460
iQ TM Supermix	1x	10	1000
MgCl ₂ (25 mM)	4.0 mM	0.8 ¹	80
BSA (20 µg µl ⁻¹)	0.2 µg µl ⁻¹	0.2	20
338F (10 µM)	0.5 µM	1.0	100
805R (10 µM)	0.5 µM	1.0	100

516P (10 µM)	0.2 µM	0.4	40
Template		2	2 x 100

1 Buffer contains MgCl₂ at final conc. of 3.0 mM

Thermocycler programme

3

1. 95 °C for 00:05:00
2. x 40 {
 - 2.1 95 °C for 00:00:30
 - 2.2 62 °C for 00:00:30 take snapshot
- }