JAN 22, 2021

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.brrmm546](https://doi.org/10.17504/protocols.io.brrmm546)

External link:
<https://doi.org/10.1371/journal.pone.0240946>

Protocol Citation: Nataly Ruiz Quinones, Cristian Daniel Asmat Ortega, Rene Flores Clavo 2021. 16s rDNA gene amplification protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.brrmm546>

MANUSCRIPT CITATION:
 Clavo RF, Quiñones NR, Hernández-Tasco

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jan 22, 2021

Last Modified: Jan 22, 2021

🌐 16s rDNA gene amplification protocol

Nataly Ruiz Quinones¹, Cristian Daniel Asmat Ortega²,
 Rene Flores Clavo¹

¹Universidade Estadual de Campinas;

²Centro de Investigación en Innovación en Ciencias Activas Multidisciplinarias

Rene Flores Clavo: Centro de Investigación e Innovación en Ciencias Activas Multidisciplinarias;

RENE FLORES

ciicam

Nataly Ruiz Quinones

Universidade Estadual de Campinas, CENTRO DE INVESTIGACIÓN E...

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Keywords: PCR amplification,
universal bacterial primers,
16s rDNA

MATERIALS

Materials

Polypropylene tubes of 1,5 mL
Polypropylene tubes of 0,5 mL
Absorbent paper
Tips of 10 µL, 200 µL, 1000 µL
Permanent marker for labeling
Gloves

Reagents

Taq buffer (10X)
Taq DNA polymerase (5 U/µL)
MgCl₂ (50 mM)
dNTP (2.5 mM each dNTP)
Primers (20 µM)
Syber safe (10000X)
Loading buffer dye (6X)
DNA Ladder 1Kb (50 ng)
TBE 1X (89mM Tris-borate, 89mM boric acid, 2mM EDTA)

Solutions

DNA template (50-100 ng/µL)
Sterile deionized water

Other

Micropipette of 10 µL, 200 µL, 1000 µL
Analytical balance
Freezer
Thermocycler

Master mix preparation

- 1 In a polypropylene tube, mix the following reagents for the final volume of 25 µL per reaction
 - 1.1 Add 2.5 µL of 10X Taq buffer (final concentration 1X)

- 1.2** Add 0.75 μL of MgCl_2 (50 mM) (final concentration 2.5 mM)
- 1.3** Add 0.2 μL dNTP (2.5 mM each dNTP) (final concentration 0,02 mM)
- 1.4** Add 0.5 μL of each primer (20 μM) (final concentration 0,4 μM)
Primer 10f 5'-GAG TTT GAT CCT GGC TCA G-3'
Primer 1401r 5'-CGG TGT GTA CAA GGC CCG GGA ACG-3' or 1492r 5'- ACC TTG TTA CGA CTT-3' or
Primer 1492r 5'- ACC TTG TTA CGA CTT-3'
- 1.5** Add 5.0 μL of the DNA template (50-100 ng/ μL) (final concentration at 1-2,5 ng per reaction)
- 1.6** Add 0,2 μL of Taq DNA polymerase (5 U/ μL) (Final concentration 0,04 U)
- 1.7** Add sterile deionized water until filling at 25 μL per reaction.

PCR reaction

- 2** For PCR amplification, use the following cycling program
 - 2.1** Run an Initial denaturation step of 5 min at 95°C

- 2.2 Run 30 cycles of 1 min at 94°C for denaturation, 1 min at 55°C for annealing, 3 min at 72°C for extension
- 2.3 Run a 3 min at 72°C for the final extension
- 2.4 Store amplification products at 4°C until visualization.

Visualization

- 3 To analyze and visualize the amplified products, put 2 µL of the PCR product with 2 µL of loading buffer dye in agarose gel electrophoresis with 1X TBE stained with SYBER safe at 1:10000.
- 4 To compare, add 1 µL of 1kb DNA ladder (final concentration 0,5 ng/µL) and observe a band of 1500 pb approximately.