16s rDNA sequencing reaction and precipitation protocol

Rene Flores Clavo¹, Cristian Daniel Asmat Ortega²,
Nataly Ruiz Quinones¹

¹Universidade Estadual de Campinas;

²Centro de Investigación en Innovación en Ciencias Activas
Multidisciplinarias

Rene Flores Clavo: Centro de Investigación e Innovación en Ciencias Activas
multidisciplinarias-CIICAM;

Nataly Ruiz Quinones: Centro de Investigación e Innovación en Ciencias Activas
multidisciplinarias-CIICAM

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.brzpm75n](https://doi.org/10.17504/protocols.io.brzpm75n)

External link:
<https://doi.org/10.1371/journal.pone.0240946>

Protocol Citation: Rene Flores Clavo, Cristian Daniel Asmat Ortega, Nataly Ruiz Quinones 2021. 16s rDNA sequencing reaction and precipitation protocol . **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.brzpm75n>

MANUSCRIPT CITATION:
Clavo RF, Quiñones NR, Hernández-Tasco

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 01, 2021

RENE FLORES

Tech. support phone: **+55 (19) 991640041** email: renefloresclavo@gmail.com
Click here to message tech. support

Rene Flores Clavo

Universidade Estadual de Campinas, Centro de Investigación e...

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

PROTOCOL integer ID:
46863

Keywords: 16s rDNA
sequencing, precipitation
reaction, protocol

Materials

Polypropylene tubes of 1,5 mL
Absorbent paper
Tips of 10 µL, 200 µL, 1000 µL
Permanent marker for labeling
Nitrile gloves
96-well sequencing plates

Reagents

Primer 10f, 765f, 782r, 1100r and 1492r (20 µM each)
The BigDye® Terminator v3.1
EDTA (125 mM)
Sodium Acetate (NaOAc) 3M
Etanol 100%
Etanol 70%
Hi-Di™ Formamide - Thermo Fisher Scientific

Solutions

Save money buffer:

- 2 ml of tris HCl pH9.0, 1M
- 1 ml of MgCl 250 mM
- Fill with deionized sterile water up to a volume of 10 ml

Tris-HCl pH9.0, 1M:

- 1,211 g of Tris and fill with deionized sterile water up to a volume of 8 ml
- Adjust pH by adding HCl 100%
- fill with deionized sterile water up to a volume of 10 ml

DNA template (10 - 20 ng/µL)
Sterile deionized water

Other

Micropipette of 10 µL, 200 µL, 1000 µL
Analytical balance
Freezer
Thermocycler
Refrigerated centrifuge for sequencing plates
Genetic Analyzer ABI 3500 series Applied Biosystems

Sequencing reaction

1

Consider a total volume of 10 μ l

2 Add 6 μ l of water for PCR

3 Add 2 μ l of save money buffer

4 Add 0.5 μ l of primer

5 Add 0.5 μ l of Byg Dye

6 Add 1 μ l of DNA (10 - 20 ng/ μ l)

7 Place the reaction in the thermal cycler with the following cycling program:

7.1 Run an Initial denaturation step of 1 min at 96 °C.

7.2 Run 30 cycles of 15 sec at 96 °C for denaturation, 15 sec at 50 °C for annealing, 4 min at 60 °C for extension.

7.3 Store amplification products at 4°C until further use in precipitation reactions.

Precipitation

- 8** Add 1 µl of EDTA (125 mM) to each well
- 9** Add 1 µl of 3M Sodium Acetate (NaOAc) to each well
- 10** Add 25 µl of 100% cold Ethanol to each well, cap tightly and vortex slightly
- 11** Incubate at 4 °C for 15 minutes
- 12** Centrifuge at 2250g for 30 minutes
- 13** Dispense the contents in the sink immediately and invert immediately on paper towels
- 14** If it is not dispensed immediately, centrifuge at 2250g for 2 min

- 15 Additional 35 µl of 70% chilled ethanol in each well
- 16 Centrifuge at 2250g for 15 minutes
- 17 Dispense the contents in the sink and centrifuge the inverted plate 15g for 1 minute
- 18 Incubate the uncapped plate in the thermal cycler at 95 °C for 5 minutes
- 19 Resuspend the samples in 10 µl of Hi-Di formamide, cover with the cap and vortex slightly
- 20 Incubate in the thermal cycler at 95 °C for 2 minutes
- 21 Put ice for 2 minutes
- 22 Place the card in the sequencer