JAN 14, 2020

2ndline_sh_and_VL

PLOS One

Rahel Bircher¹

¹Schweizerisches Tropen- und Public Health Institut

Rahel Bircher

ATTACHMENTS

[SOP_2ndline_VL.pdf](#) [SOP_2ndline_sh.pdf](#)

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.749hqz6

External link:

<https://doi.org/10.1371/journal.pone.0227600>

Protocol Citation: Rahel Bircher 2020.

2ndline_sh_and_VL.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.749hqz6>

MANUSCRIPT CITATION:

Bircher RE, Ntamatungiro AJ, Glass TR, Mnzava D, Nyuri A, Mapesi H, Paris DH, Battegay M, Klimkait T, Weisser M, obotKsg (2020) High failure rates of protease inhibitor-based antiretroviral treatment in rural Tanzania – A prospective cohort study. PLoS ONE 15(1): e0227600. doi:

[10.1371/journal.pone.0227600](https://doi.org/10.1371/journal.pone.0227600)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

ATTACHMENTS

[SOP_2ndline_VL.pdf](#)

[SOP_2ndline_sh.pdf](#)

ATTACHMENTS

[SOP_2ndline_VL.pdf](#)

[SOP_2ndline_sh.pdf](#)

Protocol status: Working

Created: Oct 11, 2019

Last Modified: Jan 14, 2020

PROTOCOL integer ID:
28545