OCT 26, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bnzdmf26

Protocol Citation: Seth Currlin 2020. 3D Immunostaining for CLARITY-processed samples. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bnzdmf26>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 26, 2020

Last Modified: Oct 26, 2020

PROTOCOL integer ID:
 43781

3D Immunostaining for CLARITY-processed samples

Seth Currlin¹

¹University of Florida

Human BioMolecular Atlas Program (HuBMAP) Method Development Community
 Tech. support email: Jeff.spraggins@vanderbilt.edu

Seth Currlin

ABSTRACT

This is a guide for immunostaining CLARITY-processed samples.
dx.doi.org/10.17504/protocols.io.8jihuke

These steps are meant to be a guide for immunostaining large samples and should be optimized to suit your particular tissues and reagents. Large tissue volumes and dense tissue types will require longer incubation and wash times. The parameters suggested below are for a piece of human thymus tissue approximately 5 mm³ in size.

A useful link: <http://wiki.claritytechniques.org/index.php/Immunostaining>

MATERIALS

Solutions:

Wash buffer (PBST): 1x phosphate buffered saline (PBS), 0.5% Triton X-100, pH 7.4; at least 20x volume of tissue.

Blocking buffer: 1x PBS, 1% horse serum, 0.5% Triton-X 100, 0.05% Tween-20, pH 7.4; at least 10x volume of tissue.

oAnother serum should be used based on the secondary antibody.

Staining buffer: 1x PBS, 0.5% Triton X-100, 0.05% Tween-20, pH 7.4; at least 10x volume of tissue.

BEFORE START INSTRUCTIONS

Ensure the tissue is sufficiently transparent for your imaging requirements. If the sample has not been successfully cleared then subsequent imaging will be difficult due to regions retaining light-obscuring lipids. Samples should be imaged prior to immunostaining to confirm adequate clearing, particularly among excitation/emission settings intended for later use.

Immunohistochemistry for large (5 mm³) tissue volumes 3w 5d

- 1** Wash out residual clearing solution 3d

Three washes in PBST, 24 hours each, 37C with gentle agitation.
- 2** Nonspecific Epitope Blocking 3d

Blocking buffer incubation: three days, 37C with gentle agitation.

Note: Serum type should be used based on the secondary antibody host species.
- 3** Primary Antibody Incubation 1w

1:100 dilution (~10 ug/mL) in Staining Buffer.

Primary antibody incubation: five days, 37C with gentle agitation; then two days, 4C with gentle agitation.
- 4** Wash out Primary Antibody 3d

Three washes, 24 hours each, at 37C with gentle agitation.
- 5** Secondary Antibody Incubation 1w

1:200 dilution (~5 ug/mL) in Staining Buffer.

Secondary antibody incubation: five days at 37C with gentle agitation; two days at 4C with gentle agitation.

Note: Protect samples from light.
- 6** Wash out Secondary Antibody 3d

Three washes, 24 hours each, at 37C with gentle agitation.

Note: Protect samples from light.

7 Prepare for 3D Imaging or Storage

Prepare for imaging or store samples in washing buffer (PBST) at 4C.

For 3D imaging guide, see: <https://dx.doi.org/10.17504/protocols.io.begajbse>