


Jan 06, 2020 Version 1

## 4-methylumbelliferyl heptanoate (MUH) - Cell Viability Assay V.1

DOI

[dx.doi.org/10.17504/protocols.io.bazgif3w](https://dx.doi.org/10.17504/protocols.io.bazgif3w)



Peter Vangheluwe<sup>1</sup>, Shaun Martin<sup>1</sup>, Mujahid Azfar<sup>1</sup>

<sup>1</sup>KU Leuven



Peter Vangheluwe

KU Leuven

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.bazgif3w](https://dx.doi.org/10.17504/protocols.io.bazgif3w)

**Protocol Citation:** Peter Vangheluwe, Shaun Martin, Mujahid Azfar 2020. 4-methylumbelliferyl heptanoate (MUH) - Cell Viability Assay. [protocols.io](https://dx.doi.org/10.17504/protocols.io.bazgif3w) <https://dx.doi.org/10.17504/protocols.io.bazgif3w>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**Wrong file. Please go to Version 2.0 of this protocol**

**Created:** January 06, 2020

**Last Modified:** May 31, 2024

**Protocol Integer ID:** 31496

**Keywords:** Cell Viability Assay, MUH assay, 4-methylumbelliferyl heptanoate, cell-mediated cytotoxicity, cell proliferation, ASAPCRN



## Abstract

The 4-methylumbelliferyl heptanoate (*MUH*) assay is a A fluorimetric assay used to study cell-mediated cytotoxicity. The principle behind this assay employs the activity of esterases found in viable cells. These esterases then hydrolyse the fluorochrome in the MUH reagent which can then be read using a plate reader.

## Attachments



Cell fractionation.p...

583KB

