

Feb 02, 2016

Marine Purity Broth

DOI

dx.doi.org/10.17504/protocols.io.dcm2u5

Bertillson et al.¹

¹Matthew Sullivan Lab, University of Arizona, Ohio State University

VERVE Net

Sullivan Lab



Verve Team

University of Arizona

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.dcm2u5

Protocol Citation: Bertillson et al. 2016. Marine Purity Broth. [protocols.io https://dx.doi.org/10.17504/protocols.io.dcm2u5](https://dx.doi.org/10.17504/protocols.io.dcm2u5)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: July 10, 2015

Last Modified: November 25, 2017

Protocol Integer ID: 1133

Abstract

Purpose: To test axenic cultures for purity

Guidelines

Formulation:

17 g	BD Difco AC Broth
20 g	NaCl
Qs to 1 L with distilled H ₂ O	

After broth has cooled:

2 M MgSO ₄	(2 M MgSO ₄ ·7H ₂ O = 49.3 g/100 ml H ₂ O, dissolve and 0.2 µm filter)
1 M CaCl ₂	(1 M CaCl ₂ ·2H ₂ O = 14.7 g/100 ml H ₂ O, dissolve and 0.2 µm filter)

Bertillson et al. 2003 Limnol Oceanogr 48(5): 1721-1731

- 1 Autoclave BD Difco AC Broth and NaCl Qs to 1 L with distilled water for 20 min.

⌚ 00:20:00

Note

See formulation in guidelines.

- 2 After broth has cooled, add 16 ml 2 M MgSO₄ and 10 ml 1 M CaCl₂

Note

(2 M MgSO₄·7H₂O = 49.3 g/100 ml H₂O, dissolve and 0.2 µm filter)

Note

(1 M CaCl₂·2H₂O = 14.7 g/100 ml H₂O, dissolve and 0.2 µm filter)

- 3 Dispense 2.0 or 4.5 ml per sterile tube

- 4 Inoculate sterile MPB with 1:5 (vol:vol) dilution of culture

- 5 Incubate at 22°C

Note

Heterotrophs normally grow in less than 2 days. No growth in the broth after 1-2 weeks indicates culture is axenic.