

Jul 12, 2016

OmniPrep™ For High Quality Genomic DNA Extraction From Nucleated Blood Cells (Bird, Fish, Frog)

 In 1 collection

DOI

dx.doi.org/10.17504/protocols.io.e54bg8w

G-Biosciences

G-Biosciences



Colin Heath

G-Biosciences



Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.e54bg8w>

External link: https://www.gbiosciences.com/image/pdfs/protocol/786-136_protocol.pdf

Protocol Citation: G-Biosciences 2016. OmniPrep™ For High Quality Genomic DNA Extraction From Nucleated Blood Cells (Bird, Fish, Frog). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.e54bg8w>



License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: June 16, 2016

Last Modified: February 15, 2018

Protocol Integer ID: 2972

Keywords: high quality genomic dna extraction, dna preparation, omniprep, genomic dna, high quality genomic dna, high quality genomic dna from many different species, dna, use with nucleated blood cell, nucleated blood cell, fish

Abstract

The OmniPrep™ kit isolates high quality genomic DNA from many different species and tissue types including animal, plant, bacteria, yeast, fungi, whole blood, and cells in culture. DNA can be isolated from samples high in polysaccharides or other contaminants that are difficult to remove from the DNA preparations.

This protocol is for use with **nucleated blood cells** (bird, fish, frog). Please refer to the **appropriate protocol** depending on your application.

Guidelines

INTRODUCTION

The OmniPrep™ kit isolates high quality genomic DNA from many different species and tissue types including animal, plant, bacteria, yeast, fungi, whole blood, and cells in culture. DNA can be isolated from samples high in polysaccharides or other contaminants that are difficult to remove from the DNA preparations.

Several unique features separate the OmniPrep™ kit from other methods:

- A unique and proprietary formulation of detergents and salts. DNA can be extracted and purified from almost any tissue without the use of toxic agents such as phenol.
- A quick protocol, which isolates extremely clean genomic DNA. On an average, the DNA is 100kb in size and generally hydrates in minutes.
- Provides an option to modify protocol for difficult to handle samples.
- Suitable for 200 Preps (10mg/prep). The kit is adaptable for larger tissue volumes.

ITEM(S) SUPPLIED

	Description	Cat. # 786-136	Cat. # 786-136S
	Genomic Lysis Buffer	100ml	2 × 2ml
	Nuclei Isolation Buffer	2 × 30ml	-
	DNA Stripping Solution	10ml	0.5ml
	Precipitation Solution	30ml	2ml
	Muscle Glycogen (10mg/ml)	1ml	25µl
	TE Buffer	20ml	0.5ml
	Longlife™ RNase (5mg/ml; >60U/mg)	0.5ml	50µl
	Longlife™ Proteinase K (5mg/ml)	2 × 0.5ml	50µl

NOTE: Cat. # 786-136S is a trial/sample size and does not contain enough reagents for all the protocols. For regular size kit, order Cat. # 786-136.

STORAGE CONDITIONS

The kit is shipped at ambient temperature. Upon arrival, store the kit components as recommended on the reagent label.

ADDITIONAL REAGENTS REQUIRED

- For all samples: Isopropanol, 70% Ethanol, and Chloroform
- For Gram positive Bacteria: Lysozyme, 0.5M EDTA. Longlife™ Lysozyme, (Cat# 786-037) available.
- For Yeast: Zymolyase®, β-mercaptoethanol, phosphate buffered saline (PBS). Longlife™ Zymolyase® (Cat# 786-036) available.

PREPARATION BEFORE USE

Proteinase K Solution: To avoid repeated freezing-thaw, dispense the Proteinase K solution into aliquots of 30µl/tube and freeze at -20°C.

Genomic Lysis Buffer & DNA Stripping Solution: If a precipitate forms due to cold storage allow to warm to room temperature until precipitate dissolves.

PROTOCOLS

The protocols contained in this manual are listed below and are based on the Extraction from Solid Tissue protocol. Thoroughly review the appropriate protocols before commencing isolation of genomic DNA.

TROUBLESHOOTING

• POOR DNA RECOVERY:

- o Increase volumes of Genomic Lysis Buffer, DNA Stripping Solution and Precipitation Solution proportionally.
- o Introduce optimal Proteinase K step after step 4.
- o Improve grinding technique.
- For efficient grinding of small samples we offer Molecular Grinding Resin™ (Cat. # 786-138), high tensile micro-particles that do not bind nucleic acids and are recommended for grinding and isolation of genomic DNA.
- DNA/RNA free matching pestles and microfuge tubes (1.5ml) for grinding small samples are also available (Cat. # 786-138P).

• FOR >100kb GENOMIC DNA

- o Recommend using MegaLong™ DNA isolation kit (Cat. # 786-146, 786-147). Isolates nuclei under mild condition, which are then transferred to Tube-ODIALYZER™, a new device for DNA isolation. Nuclei are digested with protease followed by dialysis to remove protein and other contaminants. For more information call our Technical Department.

Typical Yield: 2-3µg DNA /10⁶ Cells

CITATIONS

1. Staley, C.A. et al (2012) Gene. 496:118
2. Jonkers, W. et al (2012) Appl Envir Microbiol 78:3656
3. Li, Z. et al (2010) Protein Expression and Purification. 72:113
4. Li, Z. et al (2010) Biochem and Biophys Res Comm. 402:519
5. Lorch, J. et al (2010) J Vet Diagn Invest 22:224



6. Li, Z. et al (2009) Protein Expression and Purification. 67:175
7. Lin-Cereghino, J. et al (2008) Yeast. 25:293
8. Choi, Y. and Shim, W. (2008) Microbiology. 154: 326
9. Choi, Y. et al (2008) Mycologia. 100:701
10. Whitaker, V. et al (2007) J. Amer. Soc. Hort. Sci. 132:534
11. Szabo, L. J. (2007) Mol Ecol Notes 7:92
12. Ordóñez, M.E. and Kolmer, J.A. (2007) Phytopathology 97:574
13. Sagaram, U.S. et al (2006) Mol Plant Pathol 7:381
14. Mertens, J.A. et al (2006) Arch. Microbiol. 186:41
15. Lee, B et al (2005) Genome. 48:1104
16. Pliss, L. et al (2004) J. Neurochem. 91:1082
17. Jacobs-Helber, S. et al (2002) JBC. 277:4859
18. Li, X. et al (2002) Genome. 45:229
19. Villar, M. et al (2001) J. Bact. 183:55

Materials

MATERIALS

 OmniPrep™ G-Biosciences Catalog #786-136

Troubleshooting

Before start

Proteinase K Solution: To avoid repeated freezing-thaw, dispense the Proteinase K solution into aliquots of 30µl/tube and freeze at -20°C.

Genomic Lysis Buffer & DNA Stripping Solution: If a precipitate forms due to cold storage allow to warm to room temperature until precipitate dissolves.




- 1 Add 10µl nucleated blood to a 1.5ml microfuge tube containing 500µl Lysis Buffer.


Note

NOTE: For larger volume of nucleated blood, use a 15ml tube and add 1ml Lysis Buffer for each 10µl blood sample. Adjust other reagents accordingly, 1ul nucleated blood is equivalent to 1mg tissue in the table in the Guidelines.

- 2 Incubate the sample at 55-60°C for 15 minutes. Do not heat higher than 60°C.

 00:15:00


- 3 **OPTIONAL:** For maximum DNA recovery, add 1µl Proteinase K solution for every 100µl Lysis Buffer and incubate at 60°C for 1-2 hours. Invert the tube periodically each hour. This step will digest hard to handle tissues and significantly improve the yield.

 00:01:00

- 4 Allow the sample to cool to room temperature.


- 5 Add 200µl chloroform and mix by inverting the tube several times.

- 6 Centrifuge for 10 minutes at 14,000xg and carefully remove the upper phase to a clean microcentrifuge tube.

 00:10:00

- 7 Add 50µl DNA Stripping Solution to the sample and invert several times to mix.

- 8 Incubate the sample for 5-10 minutes at 60°C.


 00:05:00

- 9 Add 100µl Precipitation Solution and mix by inverting the tube several times.



Note

A white precipitate should be produced, if not add 50µl aliquots of Precipitation Solution until a white precipitate forms.

- 10 Centrifuge the sample at 14,000xg for 5 minutes.

 00:05:00



- 11 Transfer the supernatant to a clean tube and precipitate the genomic DNA with 500µl isopropanol.
- 12 Invert the tubes 10 times to precipitate the DNA.
- 13 **OPTIONAL:** For increased DNA recovery, add 2µl Mussel Glycogen as a DNA carrier.
- 14 Centrifuge at 14,000xg for 5 minutes to pellet genomic DNA.
 00:05:00
- 15 Remove the supernatant.
- 16 Add 700µl 70% ethanol to the tube and invert several times to wash the DNA pellet.
- 17 Centrifuge for 1 minute at 14,000xg.
 00:01:00
- Note

In some samples, the pellet may be hard to see at this point and will be loosely attached to the tube.
- 18 Decant or pipette off the ethanol wash.
- 19 Invert the tube on a clean absorbent surface for several minutes to allow any excess ethanol to drain away.

Note

Do not let the pellet dry completely or it will be difficult to rehydrate.
- 20 Add 50µl TE Buffer to the pellet.
- 21 Incubate at room temperature for at least 15 minutes to rehydrate.



00:15:00

Note

Incubating the tube at 55-60°C will speed up rehydration. Incubate for 5-60 minutes.

22 **OPTIONAL:** 1µl LongLife™ RNase for every 100µl TE Buffer can be added at this stage.

23 Store DNA at 4°C, for long term storage store at -20°C or -80°C.

Note

Typical Yield: 2-3µg DNA /10⁶ Cells