

Mar 04, 2016

## TE Buffer

 Evolutionary Applications

DOI

[dx.doi.org/10.17504/protocols.io.dyw7xd](https://dx.doi.org/10.17504/protocols.io.dyw7xd)

Andrew Baker<sup>1</sup>, Ross Cunning<sup>1</sup>

<sup>1</sup>University of Miami



Andrew Baker

### Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.dyw7xd>

External link: <https://doi.org/10.1111/eva.12408>

**Protocol Citation:** Andrew Baker, Ross Cunning 2016. TE Buffer. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.dyw7xd>

**Manuscript citation:**

Putnam HM, Davidson JM, Gates RD, Ocean acidification influences host methylation and phenotypic plasticity in environmentally susceptible corals. *Evolutionary Applications* 9(9). doi: [10.1111/eva.12408](https://doi.org/10.1111/eva.12408)

**License:** This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**Created:** October 11, 2015

**Last Modified:** November 09, 2017

**Protocol Integer ID:** 1782

**Keywords:** te buffer so that edta, te buffer this te buffer, te buffer, concentration of edta, taq polymerase, buffer, taq polymerase in downstream pcr reaction, mm edta for storage, nucleic acid, edta, downstream pcr reaction, mm edta, pcr

## Abstract

This TE buffer contains 10 mM Tris and 0.1 mM EDTA for storage of nucleic acids. Concentration of EDTA is lower than other formulations of TE buffer so that EDTA will not inhibit Taq Polymerase in downstream PCR reactions.

## Troubleshooting

