MAY 23, 2018

A General Protocol for Western Blotting Mammalian Cell Lysates

David Dilworth¹

¹Structural Genomics Consortium

David Dilworth

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.pxndpme

Protocol Citation: David Dilworth 2018. A General Protocol for Western Blotting Mammalian Cell Lysates.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.pxndpme>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 06, 2018

Last Modified: May 23, 2018

PROTOCOL integer ID:
11982

Harvesting and Lysis

- 1 For adherent cell lines, wash with 1 x PBS and detach cells by incubating in 0.25% trypsin. Once cells have detached at an equal volume of complete media and transfer to a microfuge tube.

Suspension cells can be transferred directly to microfuge tube from culture. Spin cells at 500 x G for 5 min to pellet. Wash pellet with 1 x PBS and repeat centrifugation followed by removal of supernatant.

- 2 Resuspend cell pellet in 50 μ L - 100 μ L lysis buffer. Incubate on ice for 10 minutes and then add SDS to 1% final.

Note

Lysis Buffer:

- * 20 mM Tris-HCl pH8

- * 150 mM NaCl

- * 10 mM MgCl₂

- * 1mM EDTA

- * 0.5 % Triton X-100

Add fresh protease Inhibitors (100x) & benzoase (10 000x) prior to lysis.

- 3 Quantify and normalize protein concentration between samples using the bicinchoninic acid assay (BCA assay) - Pierce BCA Protein Assay Kit (Cat# 23225)

Western Blotting

- 4 To each sample add SDS Loading buffer to 1 x and boil samples for 5 min.

98 °C

- 5 Load SDS page with 50 - 100 μ g total protein and run in an appropriate buffer at 100 V for ~ 2 hours, until the dye front runs off. For NuPAGE 4-12% Bis-Tris Protein Gel (NP0322BOX), we run in 1x MOPS Running Buffer.
- 6 Transfer proteins using appropriate transfer apparatus, for 1.5 hrs at 80 volts in 1 x Tris-Glycine transfer buffer to a 0.2 μ m PVDF membrane. For PVDF membranes, pre-soak in 100% methanol followed by transfer buffer.
- 7 Block membrane in 5% milk in PBS-T (1x PBS - 0.1% Tween-20) for 30 min at RT.

- 8** Cut membrane and probe with desired antibodies diluted in 5% BSA in PBS-T overnight at 4oC.
- 9** Wash membranes 3 x - 10min in PBS-T.
- 10** Incubate membranes in secondary LiCor antibodies to mouse and/or rabbit (diluted - 1:5000) in Licor Odyssey Blocking buffer (927-40000) diluted 1 in 5 in PBS-T.
- 11** Wash membranes 3 x - 10min in PBS-T.
- 12** Image blots on Licor Odyssey CLx Imaging System.