

May 15, 2020 Version 2

## A method for isolating RNA from canine bone V.2

 In 1 collection

DOI

[dx.doi.org/10.17504/protocols.io.bf9vjr66](https://dx.doi.org/10.17504/protocols.io.bf9vjr66)



Rebecca Nance<sup>1</sup>, Payal Agarwal<sup>1</sup>, Maninder Sandey<sup>2</sup>, Dmytro Starenki<sup>3</sup>, Jey Koehler<sup>2</sup>, Abdul Mohin Sajib<sup>1</sup>, Bruce F Smith<sup>1</sup>

<sup>1</sup>Scott-Ritchey Research Center, Auburn University College of Veterinary Medicine;

<sup>2</sup>Department of Pathobiology, Auburn University College of Veterinary Medicine;

<sup>3</sup>Hudson Alpha Institute for Biotechnology, Huntsville AL



**Rebecca Nance**

University of Florida

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.bf9vjr66](https://dx.doi.org/10.17504/protocols.io.bf9vjr66)

External link: [https://www.future-science.com/doi/10.2144/btn-2019-0153?](https://www.future-science.com/doi/10.2144/btn-2019-0153?fbclid=IwAR1AdUnM10wOsts2pwp5AYhi_UkxRibNLK4ZjWFpwuusnqA6_lyRxilFeGg)

[fbclid=IwAR1AdUnM10wOsts2pwp5AYhi\\_UkxRibNLK4ZjWFpwuusnqA6\\_lyRxilFeGg](https://www.future-science.com/doi/10.2144/btn-2019-0153?fbclid=IwAR1AdUnM10wOsts2pwp5AYhi_UkxRibNLK4ZjWFpwuusnqA6_lyRxilFeGg)

**Protocol Citation:** Rebecca Nance, Payal Agarwal, Maninder Sandey, Dmytro Starenki, Jey Koehler, Abdul Mohin Sajib, Bruce F Smith 2020. A method for isolating RNA from canine bone . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bf9vjr66>

### Manuscript citation:

Nance R, Agarwal P, Sandey M, et al. A method for isolating RNA from canine bone. Biotechniques, 2020. Published online 17 Apr 2020.

**License:** This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** May 12, 2020

**Last Modified:** May 15, 2020

Protocol Integer ID: 36885

**Keywords:** bone, canine, dog, osteoblasts, osteoclasts, osteocytes, RNA, sequencing ,

## Abstract

Extracting sufficient quantity and quality RNA from bone is essential for downstream application, such as transcriptomic sequencing, to evaluate gene expression. Isolation of RNA from bone presents a unique challenge owing to the hypocellular, brittle and mineralized matrix, which makes homogenizing the tissue difficult and provides little RNA to work with. Removal of contaminating tissue, such as bone marrow and connective tissue, is essential for isolating RNA that is unique to osteoblasts, osteoclasts and osteocytes. This protocol establishes a method to effectively isolate RNA from normal canine bone cells using the second phalanx, without contamination from other tissue types, for downstream transcriptomic analysis.

This method combines physical manipulation to remove exterior tissue, washing and centrifugation to remove cells and fat within the diaphysis, homogenization using a mortar and pestle on dry ice prior to bead dissociation, followed by acid guanidinium thiocyanate-phenol-chloroform extraction and column purification to yield sufficient quantity and quality RNA from canine phalanges. The second phalanx was chosen due to its size small enough to fit into a 1.7 mL microfuge tube, but large enough to provide enough RNA. Mean RNA obtained using this protocol was 14.7 ug of RNA per gram of frozen bone.

## Attachments











[btn-2019-0153.pdf](#)

1.2MB


## Materials




### MATERIALS

-  KimWipes **Fischer Scientific**
-  TRI Reagent® **Molecular Research Center, Inc.**
-  RNase AWAY®; Spray Bottle, RNase in spray bottle; 475mL **Thermo Fisher Catalog #7002**
-  Ceria stabilized zirconium oxide beads 0.5 mm diameter **Next Advance Catalog #ZrOB05**
-  Bullet Blender **Next Advance Catalog #Bullet Blender 24**
-  Bromochloropropane (BCP) **Molecular Research Center, Inc. Catalog #BP151**
-  DNase I **Thermo Fisher Scientific Catalog #18068015**
-  RNeasy Micro Kit **Qiagen Catalog #74034**



## Before start

Wipe all surfaces and materials with RNase Away to minimize potential degradation due to RNases. Prepare steel mortar and pestle RNA-free by wiping with RNase Away, wrapping in aluminum foil, and baking at  350 °F for

 06:00:00 or  Overnight . Freeze in  -80 °C prior to RNA extraction.

## Bone Preparation


- 1 Remove all exterior soft tissue around diaphysis using scissors/scalpel/KimWipes. Use a scalpel to scrape bone along diaphysis exterior to ensure complete removal of periosteum.

### Note


\*\*Isolating RNA that is unique to osteoblasts, osteoclasts, and osteocytes is dependent on sufficient removal of these 'contaminating' exterior tissues.

- 2 Cut and discard epiphyses using large bolt cutters, being careful to not disturb the tubular structure of the diaphysis.









- 3 Place diaphysis in a 1.7 mL microfuge tube and centrifuge at  10.000 x g, Room temperature, 00:10:00 . Liquid in the bottom after centrifugation is the bone marrow/fat contained within the diaphysis.









- 4 Transfer diaphysis to a new 1.7 mL microfuge tube. Add enough 1x PBS (Phosphate Buffered Saline) to cover the bone completely. Centrifuge at




 10.000 x g, Room temperature, 00:05:00 .





- 5 Transfer diaphysis to a new 1.7 mL microfuge tube. Add enough 1x PBS to cover the bone completely and centrifuge again at  10.000 x g, Room temperature, 00:05:00 . 
- 6 Snap freeze bone in liquid nitrogen and store at  -80 °C for up to 8 months until RNA extraction. 

## Bone Homogenization/RNA Extraction

- 7 Prepare two 1.7 mL Eppendorf Safe Lock\*\* microfuge tubes, each containing  1 mL Tri-Reagent and approximately  0.5 g of zirconium oxide beads (0.5 mm diameter) and place tubes  On ice .




### Note

\*\*Eppendorf Safe Lock tubes prevent sample leakage during subsequent Bullet Blender step.



- 8 Grind bone into a fine powder using a frozen RNase-free steel mortar and pestle\*\* and hammer on top of a bed of dry ice, working quickly to avoid RNase degradation.

### Note

\*\*Mortar and Pestle can be prepared RNase-free by wiping with RNase Away, wrapping in aluminum foil, and baking at 350F for 6 hours or overnight. Freeze in  -80 °C prior to RNA extraction.






- 9 Divide the bone powder into the two previously prepared microfuge tubes containing  1 mL Tri-Reagent and  0.5 g zirconium beads .



- 10 Using a Bullet Blender at speed "5.5", subject samples to 4 rounds of a 🕒 00:00:30 spin in Bullet Blender with a 🕒 00:01:00 incubation 🌡️ On ice between rounds (to keep samples cool and minimize degradation).
- 11 Incubate samples for 🕒 00:10:00 at 🌡️ Room temperature . 📺
- 12 Add 🧪 100  $\mu$ L bromochloropropane (BCP) to each tube and vortex thoroughly. 📺 🧪  
Incubate samples for 🕒 00:05:00 at 🌡️ Room temperature .



13 Centrifuge tubes at 20.000 x g, 4°C, 00:15:00 .



- 14 Carefully remove top aqueous layer and add 10  $\mu$ L DNase I and 1/10 volume DNase I Reaction Buffer; Incubate 00:10:00 at Room temperature .



























- 15 Inactivate DNase by adding 10  $\mu$ L of 25 mM EDTA (included) and heat for 00:10:00 at 65  $^{\circ}$ C .

- 16 Add equal volume 70% ethanol and mix well by pipetting.

- 17 Apply sample to RNA column from RNeasy Micro Kit placed in a 2 mL collection tube and centrifuge 21.000 x g, Room temperature, 00:01:00 .

- 18 After discarding eluate, add 700  $\mu$ L Buffer RW1 , incubate on the column for 00:02:00 at Room temperature , centrifuge 21.000 x g, Room temperature, 00:01:00 .



- 19 Discard flow through, add  500  $\mu$ L Buffer RPE , incubate on the column for  00:02:00 at  Room temperature , and centrifuge  21.000 x g, Room temperature, 00:01:00 .   
- 20 Discard flow through, add  500  $\mu$ L of 80% ethanol , incubate on the column for  00:02:00 at  Room temperature , and centrifuge  21.000 x g, Room temperature, 00:03:00 .   
- 21 Discard flow through and replace 2 mL collection tube and centrifuge the columns at  21.000 x g, 00:05:00 , with lids open . 
- 22 Replace collection tube, add  17  $\mu$ L of RNase-free water (preheated to  65  $^{\circ}$ C ), incubate for  00:10:00 on the column at  Room temperature , and centrifuge  21.000 x g, Room temperature, 00:05:00 .   
- 23 Concentration and purity can then be determined using a Nanodrop spectrophotometer. 