

Aug 23, 2023

② A recombinant retroviral expression vector (pLXSN-HygR) based on pLXSN that confers resistance to hygromycin and pLXSN-HygR derivatives that encode EGFR, ERBB2, or ERBB3



DOI

RESEARCH

dx.doi.org/10.17504/protocols.io.j8nlkooq1v5r/v1

David J Riese II^{1,2}, Vipasha Dwivedi¹

¹Auburn University; ²University of Alabama-Birmingham



David J Riese II

Auburn University

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account





DOI: https://dx.doi.org/10.17504/protocols.io.j8nlkooq1v5r/v1

Protocol Citation: David J Riese II, Vipasha Dwivedi 2023. A recombinant retroviral expression vector (pLXSN-HygR) based on pLXSN that confers resistance to hygromycin and pLXSN-HygR derivatives that encode EGFR, ERBB2, or ERBB3. **protocols.io** https://dx.doi.org/10.17504/protocols.io.j8nlkooq1v5r/v1



License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 23, 2023

Last Modified: August 23, 2023

Protocol Integer ID: 86897

Keywords: pLXSN, hygromycin, recombinant retroviral vector, EGFR, ERBB2, ERBB3, hygromycin resistance gene, neomycin resistance gene, retroviral expression vector, resistance to hygromycin, functional effects of erbb4 heterodimerization, erbb4 heterodimerization, construction of plxsn, gene, hygr derivative, encode egfr, plxsn, erbb2, erbb3, hygr, called plxsn, egfr

Disclaimer

DISCLAIMER - FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to <u>protocols.io</u> is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with <u>protocols.io</u>, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Here we describe the construction of a derivative of pLXSN, called pLXSN-HygR, that expresses a hygromycin resistance gene rather than a neomycin resistance gene. We also describe the construction of pLXSN-HygR derivatives that express the *EGFR*, *ERBB2*, or *ERBB3* genes. Thus, these constructs are ideal for studying the functional effects of ERBB4 heterodimerization with EGFR, ERBB2, or ERBB3.

Troubleshooting



Introduction

- Recombinant retroviruses are commonly used to direct the ectopic expression of genes in infected cells. There are two significant advantages to this approach. Following infection of the target cell, the recombinant retroviral genome is reverse-transcribed and subsequently integrated into the host cell genome, enabling stable ectopic gene expression. Recombinant retroviral expression vectors typically contain a drugresistance gene, allowing the selection and maintenance of stably infected cells.
- We have previously reported the construction of recombinant retroviral expression vectors based on pLXSN [1] that express the *EGFR*, *ERBB2*, *ERBB3*, or *ERBB4* receptor tyrosine kinase genes [2, 3]. These constructs are useful for studying the role of these genes in regulating cellular proliferation, particularly in tumor cells.
- However, ERBB family receptor tyrosine kinases can undergo heterodimerization, thereby diversifying the effects of ERBB receptor signaling **[4, 5]**. Therefore, there is a need for constructs that enable the simultaneous ectopic expression of two ERBB receptor tyrosine kinase genes. Hence, here we describe the construction of a derivative of pLXSN, called pLXSN-HygR, that expresses a hygromycin resistance gene rather than a neomycin resistance gene. We also describe the construction of pLXSN-HygR derivatives that express the *EGFR*, *ERBB2*, or *ERBB3* genes. Thus, these constructs are ideal for studying the functional effects of ERBB4 heterodimerization with EGFR, ERBB2, or ERBB3.

Methods

4 Construction of pLXSN-HygR

We have previously briefly described the construction of pLXSN-HygR **[6]**. The plasmid pSV2-Hyg **[7]**, which encodes the hygromycin resistance gene **[8]**, is a generous gift from Daniel DiMaio.

- The plasmid pSV2-Hyg was digested with BgIII, after which the overhang was filled using the Klenow fragment of the *E. Coli* DNA polymerase I. This product was digested with HindIII, releasing a 1348 bp fragment that contains the hygromycin resistance gene; the hygromycin resistance gene is 1026 bp in length (nt 2050 to 3075 of pLXSN-HygR) and encodes a 341 aa protein. Therefore, this fragment was used as the insert for pLXSN-HygR. The HindIII site of this fragment lies approximately 30 nt upstream of the start codon of the hygromycin resistance gene, whereas the former BgIII site of this fragment lies approximately 290 nt downstream of the stop codon of the hygromycin resistance gene.
- The plasmid pLXSN [1, 3] (pLXSN.dna 48KB **Figure 1**) was digested with HindIII and Nael. Note that Nael produces blunt ends. The largest fragment (5071 bp) of this



digestion is missing the neomycin resistance gene and was used as the vector for pLXSN-HygR.




Figure 1. A map of pLXSN (**pLXSN.dna** – see link above) is shown. Note that the neomycin resistance gene is flanked at the 5' end by a unique HindIII site (nt 2007) and at the 3' end by a Nael site (nt 2979). Thus, the 5071 bp Nael-HindIII fragment of pLXSN serves as the vector for pLXSN-HyqR.

The pLXSN-Nael-HindIII fragment was ligated with the pSV2-Hyg-HindIII-Blunt fragment and the ligation reaction product was electro-transformed into DH10B *E. coli*. Ampicillin-resistant colonies of *E. coli* were expanded, and minipreps were screened for the presence and correct orientation of the hygromycin resistance gene insert. NGS has validated this construct, and the map of the resulting sequence (

pLXSN-HygR.dna 50KB) is shown in **Figure 2**.




Figure 2. A map of pLXSN-HygR (pLXSN-HygR.dna – see above for link) is shown. Note the unique Ascl, AvrII, XmnI, HindIII, Asel, and Stul sites used to generate pLXSN-HygR-EGFR, pLXSN-HygR-ERBB2 and pLXSN-HygR-ERBB3.

8 **Construction of pLXSN-HygR-EGFR**

To generate the pLXSN-HygR vector fragment, which contains the hygromycin resistance gene, we digested pLXSN-HygR (pLXSN-HygR.dna – see above for link) with Avrll and Ascl (Figure 2) and isolated the 5043 bp fragment. To generate the EGFR cDNA fragment, we digested pLXSN-EGFR (pLXSN-EGFR.dna 69KB) [2, 3] with Ascl and AvrII and isolated the 5521 bp fragment (Figure 3).




Figure 3. A map of pLXSN-EGFR (**pLXSN-EGFR.dna** – see above for link) is shown. Note the unique AscI site at nt 617 and the unique AvrII site at nt 6138. The 5521 bp AscI-AvrII fragment of pLXSN-EGFR encodes the *EGFR* cDNA and was used to construct pLXSN-HygR-EGFR.

The pLXSN-HygR-AvrII-AscI fragment was ligated with the pLXSN-EGFR-AscI-AvrII fragment and the ligation reaction product was electro-transformed into DH10B *E. coli*. Ampicillin-resistant colonies of *E. coli* were expanded, and minipreps were screened for the correct recombinant plasmid. NGS has validated this construct, and the map of the resulting sequence (pLXSN-HygR-EGFR.dna 68KB) is shown in **Figure 4**.




Figure 4. A map of pLXSN-HygR-EGFR (**pLXSN-HygR-EGFR.dna** – see above for link) is shown.

10 **Construction of pLXSN-HygR-ERBB2**

We have previously briefly described the construction of pLXSN-HygR-ERBB2 [9]. To generate the pLXSN-HygR vector fragment, which contains the hygromycin resistance gene, we digested pLXSN-HygR (**pLXSN-HygR.dna** – see above for link) with HindIII and XmnI (**Figure 2**) and isolated the 4017 bp fragment. To generate the *ERBB2* cDNA

fragment, we digested pLXSN-ERBB2 (pLXSN-ERBB2.dna 83KB) [2, 3] with Xmnl and HindIII and isolated the 6469 bp fragment (**Figure 5**).




Figure 5. A map of pLXSN-ERBB2 (**pLXSN-ERBB2.dna** – see above for link) is shown. Note the unique HindIII site at nt 6076 and the unique XmnI site at nt 9719. The 6469 bp XmnI-HindIII fragment of pLXSN-ERBB2 encodes the *ERBB2* cDNA and was used to construct pLXSN-HygR-ERBB2.

The pLXSN-HygR-HindIII-XmnI fragment was ligated with the pLXSN-ERBB2-XmnI-HindIII fragment and the ligation reaction product was electro-transformed into DH10B *E. coli*. Ampicillin-resistant colonies of *E. coli* were expanded, and minipreps were screened for the correct recombinant plasmid. NGS has validated this construct, and the map of the resulting sequence (pLXSN-HygR-ERBB2.dna 70KB) is shown in **Figure 6**.






Figure 6. A map of pLXSN-HygR-ERBB2 (pLXSN-HygR-ERBB2.dna - see above for link) is shown.

12 **Construction of pLXSN-HygR-ERBB3**

We have previously briefly described the construction of pLXSN-HygR-ERBB3 [9]. To generate the pLXSN-HygR vector fragment, which contains the hygromycin resistance gene, we digested pLXSN-HygR (pLXSN-HygR.dna - see above for link) with Stul and Asel (Figure 2) and isolated the 3607 bp fragment. To generate the ERBB3 cDNA

fragment, we digested pLXSN-ERBB3 (pLXSN-ERBB3.dna 69KB) [2, 3] with Asel and Stul and isolated the 7116 bp fragment (Figure 7).






Figure 7. A map of pLXSN-ERBB3 (**pLXSN-ERBB3.dna** – see above for link) is shown. Note the unique Stul site at nt 6296 and the unique Asel site at nt 9529. The 7116 bp Asel-Stul fragment of pLXSN-ERBB3 encodes the *ERBB3* cDNA and was used to construct pLXSN-HygR-ERBB3.

The pLXSN-HygR-Stul-Asel fragment was ligated with the pLXSN-ERBB3-Asel-Stul fragment and the ligation reaction product was electro-transformed into DH10B *E. coli*. Ampicillin-resistant colonies of *E. coli* were expanded, and minipreps were screened for the correct recombinant plasmid. NGS has validated this construct, and the map of the resulting sequence (pLXSN-HygR-ERBB3.dna 90KB) is shown in **Figure 8**.




Figure 8. A map of pLXSN-HygR-ERBB3 (**pLXSN-HygR-ERBB3.dna** – see above for link) is shown.

Conclusion

Here we describe the construction of a recombinant retroviral vector (pLXSN-HygR) that carries the hygromycin resistance gene instead of the neomycin resistance gene, as well as derivatives that also carry the *EGFR*, *ERBB2*, or *ERBB3* cDNAs (pLXSN-HygR-EGFR, pLXSN-HygR-ERBB2, and pLXSN-HygR-ERBB3, respectively). These constructs, together with the recombinant retroviral vector that carries the neomycin resistance gene and the *ERBB4* cDNA (pLXSN-ERBB4) [2, 3], will enable us to generate bi-recombinant cell lines to study the effects of ERBB4 receptor heterodimerization with the other ERBB family receptors.

References

15

1. Miller, A.D. and G.J. Rosman, *Improved retroviral vectors for gene transfer and expression*. Biotechniques, 1989. **7**(9): p. 980-2, 984-6, 989-90.



- 2. Riese, D.J., 2nd, et al., *The cellular response to neuregulins is governed by complex interactions of the erbB receptor family.* Mol Cell Biol, 1995. **15**(10): p. 5770-6.
- 3. Riese II, D.J. *Recombinant retroviral expression vectors based on LXSN that encode EGFR, ERBB2, ERBB3, and ERBB4*. Protocols.io 2023 [Accessed August 21, 2023]; Available from: https://dx.doi.org/10.17504/protocols.io.261gedd97v47/v1.
- 4. Riese, D.J., 2nd and D.F. Stern, *Specificity within the EGF family/ErbB receptor family signaling network.* Bioessays, 1998. **20**(1): p. 41-8.
- 5. Lucas, L.M., et al., *The Yin and Yang of ERBB4: Tumor Suppressor and Oncoprotein.* Pharmacol Rev, 2022. **74**(1): p. 18-47.
- 6. Riese, D.J., 2nd and D. DiMaio, *An intact PDGF signaling pathway is required for efficient growth transformation of mouse C127 cells by the bovine papillomavirus E5 protein.* Oncogene, 1995. **10**(7): p. 1431-9.
- 7. Lin, B.F. and B. Shane, *Expression of Escherichia coli folylpolyglutamate synthetase in the Chinese hamster ovary cell mitochondrion.* J Biol Chem, 1994. **269**(13): p. 9705-13.
- 8. Bernard, H.U., G. Krammer, and W.G. Rowekamp, *Construction of a fusion gene that confers resistance against hygromycin B to mammalian cells in culture.* Exp Cell Res, 1985. **158**(1): p. 237-43.
- 9. Feroz, K., E. Williams, and D.J. Riese, 2nd, *ErbB2 and ErbB3 do not quantitatively modulate ligand-induced ErbB4 tyrosine phosphorylation.* Cell Signal, 2002. **14**(9): p. 793-8.