AUG 17, 2020

A simple method for purification of immunoglobulin-Y (IgY).

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjuwknxe

Protocol Citation: Angel A Justiz-Vaillant 2020. A simple method for purification of immunoglobulin-Y (IgY)..

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bjuwknxe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Aug 17, 2020

Last Modified: Aug 17, 2020

PROTOCOL integer ID:

40566

1 Separate egg yolk from the egg white.

- 2 Add ten parts of distilled water to one part egg yolk.
- 3 Mix and stir slowly for 6.30 h at 4°C.
- 4 Centrifuge at 10 000 × g, at 4°C for 27 min to precipitate the lipids.
- 5 Collect the supernatant containing the IgY.
- 6 Slowly add potassium sulfate to the preparation or specimen, stirring constantly, to a final concentration of 500 mM.
- 7 Adjust pH to 7.6.
- 8 Pass the sample through a 0.45 µm filter just before applying it to the HiTrap™ Column column.