NOV 13, 2018

A Simple Method to Efficiently Record/ Capture *Caenorhabditis elegans* Locomotory Behaviours

Vishnu Raj¹, Anoopkumar Thekkuveetil¹

¹Division of Molecular Medicine, Sree Chitra Tirunal Institute for Medical Sciences and Technology

Vishnu Raj

ABSTRACT

Caenorhabditis elegans is an excellent model to study animal chemotaxis and thermotaxis behaviours. These nematodes have highly predictable behaviour pattern towards olfactory cues. A complex chemosensory information processing, based on both temporal and spacial cues, is involved in chemotaxis behaviour and can modify its behaviour towards attractants as well as repellents (Ward., 1973; Colbert et al., 1995; Troemel et al., 1997). In chemotaxis assay, worms show unsurpassed behaviour with a pattern of movement based on concentration gradient in the assay plate (Saeki et al., 2001; Iino et al., 2009). Such behavioural patterns are highly intriguing because they give better understanding on how various neuronal signalling elicit such pattern of behaviour and how factors such as past experience of the animal, mutations affecting neuronal function, modify them. (Brenner., 1974; de Bono et al., 1998). Hence, behavioural assays have critical role in elucidating the alterations in neuronal activities in *C. elegans*.

The standard chemotaxis assay estimates the movement pattern of *C. elegans* by tracking the course it takes in an agar plate containing a chemical gradient. This measurement requires recording the tracks over time in the plate (Buckingham et al., 2005; Yemini et al., 2011). Automated single worm tracker allows long term behavioural recording (Husson et al., 2005; Wang et al., 2013). The pattern of behaviour of animals in the assay plates shows there are significant alterations in patterns of movement like body bends and omega turns under experimental conditions. For a long term observation for such behaviour one needs to record the animals with least disturbances. Efficient recording often eliminates researcher's bias and makes it easy to re-evaluate the results if needed (Piere-Shimoura et al., 1999; Hardaker LA et al., 2001; Baek et al., 2002). These recorded videos can be later processed in ImageJ for measuring the patterns. Though these recording can be done using a simple dissection microscopic system with camera, there is a major limitation that the light source is very close to the worm making series of artefacts in its behaviour,

Here we report a simple setup to manually record and count these behavioural changes in worms. In this study we measured basal slowing response and enhanced slowing responses, the two different locomotory changes in response to food.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.uu9ewz6

Protocol Citation: Vishnu Raj, Anoopkumar Thekkuveetil 2018. A Simple Method to Efficiently Record/ Capture *Caenorhabditis elegans* Locomotory Behaviours. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.uu9ewz6>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Oct 20, 2018

Last Modified: Nov 13, 2018

PROTOCOL integer ID:
 17025

References

- Baek JH., Cosman P., Feng Z., Silver J., Schafer W.R. Using machine vision to analyze and classify *Caenorhabditis elegans* behavioral phenotypes quantitatively. *J. Neurosci. Meth.* 2002. 118: 9–21. PMID:12191753.
- Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics.* 1974. 77:71-94.
- Buckingham S. D., & Sattelle D. B. Fast, automated measurement of nematode swimming (thrashing) without morphometry. *B Neurosci.* 2009. 10(1):1. PMID:19619274.
- Colbert H. A., & Bargmann C. I.: Odorant-specific adaptation pathways generate olfactory plasticity in *C. elegans*. *Neuron.* 1995. 14(4):803-812. PMID:7718242.
- de Bono M., Bargmann CI. Natural variation in a neuropeptide Y receptor homolog modifies social behavior and food response in *C. elegans*. *Cell.* 1998. 94:679-89. PMID:9741632.
- Hardaker LA., Singer E., Kerr R., Zhou G., Schafer WR. Serotonin modulates locomotory behavior and coordinates egg-laying and movement in *Caenorhabditis elegans*. *J. Neuro.Bio.* 2001. 49:303-13. PMID:11745666.
- Husson S. J., Costa, W. S., Schmitt C., Gottschalk A. Keeping track of worm trackers. *WormBook.* 2005. 10.1895/wormbook.1.156.1. <http://www.wormbook.org/>
- Iino Y., & Yoshida K. Parallel use of two behavioral mechanisms for chemotaxis in *Caenorhabditis elegans*. *J. Neurosci.* 2009. 29(17):5370-5380. PMID:19403805.
- Pierce-Shimomura JT., Morse TM., Lockery SR. The fundamental role of pirouettes in *Caenorhabditis elegans* chemotaxis. *J Neurosci.* 1999. 19:9557-69. PMID:10531458.
- Saeki S., Yamamoto M., Iino Y. Plasticity of chemotaxis revealed by paired presentation of a chemoattractant and starvation in the nematode *Caenorhabditis elegans*. *J. Exp.Bio.* 2001 204(10):1757-1764. PMID:11316496.
- Sawin ER., Ranganathan R., Horvitz HR. *C. elegans* locomotory rate is modulated by the environment through a dopaminergic pathway and by experience through a serotonergic pathway. *Neuron.* 2000. 26:619–631. PMID:10896158.
- Troemel E. R., Kimmel B. E., Bargmann C. I. Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in *C. elegans*. *Cell.* 1997. 91(2):161-169. PMID:9346234.
- Wang S. J., & Wang Z. W. Track-a-worm, an open-source system for quantitative assessment of *C. elegans* locomotory and bending behavior. *Plos One.* 2013. 8(7): e69653. PMID:23922769.
- Ward S. Chemotaxis by the nematode *Caenorhabditis elegans*: identification of attractants and analysis of the response by use of mutants. *PNAS.* 1973. 70(3):817-821. PMID:4351805.
- Yemini E., Kerr R. A., Schafer W. R. Tracking movement behavior of multiple worms on food. *CSH protocols:* 2011(12):pdb-prot067025. PMID:22135669.

ATTACHMENTS

[Video-1.mp4](#) [Video-2.mp4](#)

MATERIALS

Dino-Lite microscope (Model AM4113T-GFBW), Dinocapture2.0 software (available online), Dino microscope stand (RK-10), Burette stand and clamps, Acrylic sheets (transparent, red and white), Metal stand (Steel made; Length 30cm X Breadth 15cm X Height 30cm), Manual stage-Olympus; Model No: IX2-KSP-9C13573, Philips 7W LED bulb, Computer with 2GB RAM, N2 Bristol (strain obtained from CGC, Minnesota, USA), Petri dishes (60mm), NG Media.

Setting up the camera

- The skeleton of this system is a burette stand and clamps to hold various parts (Fig.1a). For video capturing we used Dino-Lite digital microscope (Model no AM4113T-GFBW) with USB connector and its software module Dinocapture2.0 (Fig.1b) to record the data. Dino-Lite digital microscope was fixed on a table top stand (Dino-Lite RK-10) enables smooth focus adjustment and quick vertical movements (Fig.1b). An X-Y stage was placed over on a custom-made metal frame (Fig.1c). Petri dish containing the worms were placed on a transparent acrylic sheet placed on this XY stage (if the stage has a universal holder, petri dish can be directly held on the stage).
- As illumination source, 7 W led white light was used (Fig.1d). To enhance the image contrast as well as uniform illumination we used white and a red acrylic sheets (Fig.1e). The gap between LED light source and the white acrylic sheet was adjusted between 4-7 cm; adjusted based on the required intensity of light to visualize the animals clearly. The distance between white and red acrylic sheets was 7-15 cm (Fig.1e). Parallel alignment of both these sheets found to be crucial for the set up.
- The USB microscope was connected to a computer through USB cable. All recording of the video was done using the Dino-Lite software module. Using this system, we tested the animal behaviour in the presence and absence of food and could establish significant variation in body bends (Fig.1h).