SEP 19, 2019

Abeoforma whisleri transient transfection protocol

Nature Methods

Elena Casacuberta¹, Aleksandra Kozyczkowska¹, Sebastian R R Najle²

¹Institute of Evolutionary Biology; ²Center for Genomic Regulation (CRG)

Protist Research to Optimize Tools in Genetics (PROT-G)

Multicellgenomelab

[1 more workspace](#) ↓

Elena Casacuberta
Institute of Evolutionary Biology

ABSTRACT

This is a protocol for transient transfection of the Ichthyosporean Abeoforma whisleri. This protocol has an efficiency of 1-2% calculated from fluorescent positive cell from Flow Cytometry data, using the amount of cells and DNA described in the protocol. Increasing DNA from above 1 microgram/microliter does not increase transfection.

MATERIALS

MATERIALS

⊗ Buffer P3 Contributed by users Catalog #19053

⊗ PBS Contributed by users

⊗ Marine Broth 2216 BD Biosciences Catalog #279110

⊗ 4D-Nucleofector System with X Unit Lonza Catalog #AAF-1002X

⊗ A.whisleri cells Contributed by users

OPEN ACCESS

DOI:

[dx.doi.org/10.17504/protocols.io.zexf3fn](https://doi.org/10.17504/protocols.io.zexf3fn)

External link:

<https://doi.org/10.1038/s41592-020-0796-x>

Protocol Citation: Elena Casacuberta, Aleksandra Kozyczkowska, Sebastian R R Najle 2019. Abeoforma whisleri transient transfection protocol. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.zexf3fn>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 22, 2019

Last Modified: Sep 19, 2019

PROTOCOL integer ID: 21687

Keywords: LONZA, nucleofection, A.whisleri

Pre-Transfection

- Count A.whisleri cells from a culture 1 to 2 weeks old grown in Marine Broth (MB) medium at 13 degrees.

- 2 Take 2 x 10⁵ cells/per transfection condition, and spin them down at 2000g for 5 min.
- 3 Discard medium and gently resuspend cells in sterile 1xPBS and spin them down again (same conditions as above)

Transfection

- 4 Discard PBS and resuspend cells with 20-25 microliters of mixture Buffer P3 Lonza+1 to 5 micrograms of reporter plasmid+ 40 micrograms of carrier DNA (empty vector such as pUC19).
- 5 Transfer de 20-25 microlters mixture to a 16 well strips from Lonza Nucleofector (X Unit) and apply CODE EN-138.

Post-Transfection

- 6 Immediately after transfection add 80 microliters of MB directly to the well. You can transfer directly to a 6 well plate with 1ml of media, or wait for 30 minutes.
- 7 Screen for transformants after 24 hours of transfection, the number of positive cells will increase untill 48 hours.

IMPORTANT

- 8 Include always a well of cells with no DNA. A.whisleri shows autofluorescence, so this control is key to compare and decide what is real transfection and what is autofluorescence.
Use high quality DNA and as concentrated as possible. We liofilize DNA prep and resuspend them to have a concentration of at least 1.5 micrograms/microl.
Use ultrapure and concentrated carrier DNA. We use pUC19 at 20micrograms/microliter.