VERSION 1

FEB 05, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.xstfnen

Protocol Citation: Matus Valach 2019. Addition of the adaptor to RNA substrates for 3' RACE (mapping P ends).
protocols.io
<https://dx.doi.org/10.17504/protocols.io.xstfnen>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Feb 05, 2019

Last Modified: Feb 05, 2019

PROTOCOL integer ID:
 20019

Keywords: RNA, RNA-Seq, 3' RACE

🌐 Addition of the adaptor to RNA substrates for 3' RACE (mapping P ends) V.1

🔗 Forked from [Addition of the adaptor to RNA substrates for 3' RACE \(mapping OH ends\)](#)

Matus Valach¹

¹Université de Montréal, Montreal, Quebec, Canada

Matus Valach

Université de Montréal, Montreal, Quebec, Canada

ABSTRACT

Simple protocol for mapping 3'-P RNA termini by RT-PCR after the addition of a 3' adaptor using *E. coli* RtcB RNA ligase.

MATERIALS

MATERIALS

🔗 RtcB Ligase - 25 rxns **New England Biolabs Catalog #M0458S**

BEFORE START INSTRUCTIONS

Prepare samples, including controls, according to the aim of the experiment. Use the chart below to decide which enzyme is appropriate for pre-treatments of the RNA. For example, if only interested in mapping 5' RNA ends, use T4 polynucleotide kinase with ATP and without ATP to infer the original phosphorylation state of the end of interest.

Setup of the phosphorylation reaction can be found here:

[dx.doi.org/10.17504/protocols.io.cpdvi5](https://doi.org/10.17504/protocols.io.cpdvi5) However, avoid heat denaturation of the enzyme and rather purify the RNA using a trizol extraction (e.g., [http://dx.doi.org/10.17504/protocols.io.eiebcbe](https://doi.org/10.17504/protocols.io.eiebcbe)) or a column clean-up (e.g., Monarch RNA Cleanup Kits from *NEB* is optimal when interested in small RNA molecules <200 nt, which is the usual exclusion limit in other products).

Original	Original	T4PNK (+ATP)	T4PNK (+ATP)	T4PNK (-ATP)	T4PNK (-ATP)	T4PNK ⁻ 3'Pase [⊖] (+ATP)	T4PNK ⁻ 3'Pase [⊖] (+ATP)	T4PNK ⁻ 3'Pase [⊖] (-ATP)	T4PNK ⁻ 3'Pase [⊖] (-ATP)
5'	3'	5'	3'	5'	3'	5'	3'	5'	3'
P	P	P	OH	P	OH	P	P	P	P
P	OH	P	OH	P	OH	P	OH	P	OH
OH	P	P	OH	OH	OH	P	P	OH	P
OH	OH	P	OH	OH	OH	P	OH	OH	OH

1 Mix the following components (9 µL):

Component	Amount [µL]	Final concentration
RNA [1 µg]	4	100 ng/µL
5'-hydroxylated 3' RACE RNA oligo [100 µM]	0.4	
RNase-free water (ddH ₂ O)	4.6	

- *Blocking the 3' end of the 3' RACE oligo is not necessary, as 3'-OH ends are not a substrate for EcRtcB RNA ligase. 5'-end has to be hydroxylated. For practical purposes, a 5' RACE RNA oligo, which usually has both ends hydroxylated, can be used for as the 3' RACE oligo when mapping 3'-P ends by EcRtcB.*

2 Denature for 2 min at 70 °C, place on ice.

3 Mix the following components (20 µL):

Component	Amount [µL]	Final concentration
RNA + oligo mix (step 2)	9	
10× RtcB RNA ligase buffer	2	1×
50% PEG-8000	4	10%
GTP [1 mM]	2	0.1 mM
MnCl ₂ [10 mM]	2	1 mM
EcRtcB RNA ligase [15 µM]	1	0.75 µM

4 Incubate for 120 min at 37 °C.

5 Purify the RNA from the RNA ligase reaction (e.g., trizol extraction or column clean-up).

6 Proceed to RT-PCR. For the RT reaction, use 250-500 ng of the purified adaptor-ligated RNA and a primer reverse complementary to the 3' adaptor. For the PCR, use an upstream (forward) primer binding to the RNA of interest and a downstream (reverse) primer binding to the 3' adaptor.