DEC 11, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.zk4f4yw

External link:
<https://doi.org/10.7717/peerj.8157>

Protocol Citation: Kiichi Hirota, Yoshiyuki Matsuo 2019. ADP assay (Colorimetric). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.zk4f4yw>

MANUSCRIPT CITATION: Kusunoki M, Hayashi M, Shoji T, Uba T, Tanaka H, Sumi C, Matsuo Y, Hirota K, Propofol inhibits stomatoxin-1-sensitive voltage-dependent K channels in pancreatic β -cells and enhances insulin secretion. PeerJ doi: [10.7717/peerj.8157](https://doi.org/10.7717/peerj.8157)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Mar 29, 2019

ADP assay (Colorimetric)

PeerJ

Kiichi Hirota¹, Yoshiyuki Matsuo¹

¹Kansai Medical University

Yoshiyuki Matsuo

MATERIALS

MATERIALS

ADP Assay Kit (Colorimetric/Fluorometric) **Abcam Catalog #ab83359**

Standard preparation

- 1 Reconstitute the ADP standard (1 μmol) in 100 μl of water to generate a 10 mM ADP stock solution.
- 2 Dilute 10 μl of the 10 mM ADP stock solution with 90 μl of ADP Assay Buffer to prepare a 1 mM ADP standard solution.
- 3 Prepare dilutions of a ADP standard using ADP Assay Buffer (0, 0.04, 0.08, 0.12, 0.16, 0.2 mM). 100 μl of each dilution is required to set up duplicate reading (2 x 50 μl).

Sample preparation

- 4 Prepare cells in a 12-well plate.
- 5 Remove media and wash cells with cold PBS.
- 6 Lyse cells in 100 μl of ADP Assay Buffer.
- 7 Incubate the lysate for 10 min on ice.

- 8 Centrifuge at 18,000 x *g* for 5 minutes at 4°C.
- 9 Collect the supernatant in a new microcentrifuge tube and keep on ice.

Assay procedure (for the colorimetric assay)

- 10 Test all samples with a minimum of two replicates.
- 11 Dilute the lysate 3-fold in ADP Assay Buffer.
- 12 Prepare 50 µl of Reaction Mix for each reaction. Prepare an amount of Reaction Mix sufficient for all samples including ADP standards.

Component	Volume (µl)
ADP Assay Buffer	44
ADP Probe	2
ADP Converter	2
ADP Developer	2

- 13 Add 50 µl of ADP standards or samples per well to a clear 96-well plate.

- 14** Add 50 μ l of Reaction Mix to each well.
- 15** Incubate for 30 min at room temperature.
- 16** Measure the absorbance at 750 nm using iMark™ Microplate Absorbance Reader.