VERSION 2

JUN 19, 2018

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.q5ddy26

Protocol Citation: Andrew Potter 2018. Adult mouse kidney dissociation.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.q5ddy26>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 19, 2018

Last Modified: Jun 19, 2018

PROTOCOL integer ID:
13189

Adult mouse kidney dissociation V.2

Andrew Potter¹

¹Cincinnati Children's Hospital Medical Center

Human Cell Atlas Method Development Community

Andrew Potter

ABSTRACT

Protocol for adult (8-10 week) mouse kidney dissociation.

ATTACHMENTS

[kidney_12_19_17.pdf](#)

GUIDELINES

Storage Conditions of Reagents

Reagent	Storage Condition
DPBS (no Ca, no Mg)	4°C
0.5 M EDTA	room temp.
RBC Lysis Buffer	4°C
Protease from <i>Bacillus Licheniformis</i>	Store 100 µL aliquots (100 mg/mL) in DPBS at -80°C
DNase	Store 10 µL aliquots (250 U/10 µL) in DPBS at -80°C

Required Equipment

Equipment	Supplier	Catalog no.
Thermomixer C or R	Eppendorf	5382000015 / Z605271

The protocol workflow is as follows:

- A. Isolate Kidney
- B. First layer
- C. Second layer
- D. Third layer
- F. Preparing cells for Chromium

MATERIALS

MATERIALS

- ⊗ DPBS (no Ca, no Mg) Thermofisher Catalog #14190144
- ⊗ 0.5 M EDTA Ambion Catalog #AM9260G
- ⊗ RBC Lysis Buffer Sigma Catalog #R7757
- ⊗ Protease from Bacillus Licheniformis Sigma Catalog #P5380
- ⊗ DNase AppliChem Catalog #A3778
- ⊗ Thermomixer C or R Eppendorf Catalog #5382000015 / Z605271

BEFORE START INSTRUCTIONS

Prepare *Bacillus Licheniformis* enzyme mix just prior to starting dissociation:

Volume (µl)	Reagent	Final concentration
894	DPBS	1X
1	0.5 M EDTA	0.5 mM
5	DNAse 1 (250 U/10 µL)	125 U / mL
100	<i>B. Lich</i> (100 mg/mL)	10 mg/mL

+25 mg tissue / 1 mL enzyme mix

Isolate Kidney

- 1 Quickly dissect and isolate kidney and transfer to ice-cold PBS.
- 2 Remove fatty tissue and kidney capsule in ice-cold PBS.
- 3 Mince whole kidney on petri dish, on ice for 2 min until fine.
00:02:00 Mincing
- 4 On petri dish weigh out 25 mg tissue. Using razor blade, transfer tissue to 1.5 mL tube containing 1 mL of enzyme mix (10 mg/mL *B. lich*).

First layer

- 5 Incubate tube on ice for 2 min. Shake every 30 seconds.

00:02:00 Incubation on ice

00:00:30 shake every 30 sec

- 6 After 2 min total time, triturate gently 20x using 1 mL pipet set to 700 μ L.

- 7 Triturate 10x every 2 minutes for 10 additional minutes (12 min total time) while incubating on ice.

00:10:00 incubate on ice

00:02:00 Triturate 10x

- 8 Spin digest mix at 4° C 10 sec at 50 g to spin down cell clumps and leave dissociated cells in supernatant.

4 °C

00:00:10 spin 50 g for 10 sec.

- 9 Remove 80% of supernatant containing single cells and apply to 30 μ M filter on 50 mL conical; rinse filter with 8 mL ice-cold PBS/BSA 0.04% into 50 mL conical. Save conical with filter for subsequent steps.

8 mL ice-cold PBS/BSA 0.04%

Second layer

- 10 Add additional 1 mL enzyme mix (10 mg/mL b. lich) to residual tissue chunks.

1 mL b. lich enzyme mix

- 11 Triturate 10x with 1 mL pipet set to 700 μ L.

- 12** Continue digesting while shaking in thermomixer, set to 4 °C at 1200 RPM for 12 additional min (24 min total). Every 4 min passage 8X with 18 gauge needle (3X total).

00:12:00 digest in thermomixer at 1200 RPM

00:04:00 every 4 min passage 8X w/18 gauge needle (3X total)

4 °C set thermomixer to 4 °C (can leave in cold room)

- 13** Spin at 4° C 10 sec for 50 g to spin down clumps of tissue, leaving released cells in supernatant.

4 °C

00:00:10 Spin at 50 g for 10 sec.

- 14** Pipet ~80% of supernatant containing released cells to the 30 µM filter (the same tube/filter as used in previous steps). Rinse filter with 8 mL ice-cold PBS/BSA 0.04%.

8 mL ice-cold PBS/BSA 0.04%

Third layer

- 15** Add additional 1 mL enzyme mix (10 mg/mL b. lich) to residual tissue chunks.

1 mL b. lich enzyme mix

- 16** Continue dissociating remaining clumps at 1400 RPM in thermomixer at 4° C for 12 additional minutes (36 min total). Every 4 min passage 8X w/18 gauge needle w/1 mL syringe (3X total).

4 °C

00:12:00 thermomixer at 1400 RPM

00:04:00 every 4 min passage 8X with 18 gauge needle

- 17** Triturate 10x and apply total volume of remaining digest mix to the same 30-µM filter used in previous steps.

- 18** Rinse filter with 8 mL ice-cold PBS/BSA 0.04%.

8 mL ice-cold PBS/BSA
0.04%

Preparing cells for Chromium

19 Transfer flow-through to two 15 mL conicals.

20 Spin 500 G for 5 min at 4° C.

4 °C
Spinning

00:05:00 Spin at 500
g

21 Remove supernatant.

22 Re-suspend both tubes (combined) in 100 µL total volume PBS/BSA 0.04% and add 900 µL RBC lysis buffer (in 15 mL conical).

100 µL PBS/BSA
0.04%

900 µL RBC lysis
buffer

23 Triturate 20x.

24 Let sit 2 min on ice.

00:02:00 On
ice

25 Add additional 9 mL ice-cold PBS/BSA 0.04%.

9 mL ice-cold PBS/BSA
0.04%

26 Spin 500 G for 5 min at 4° C.

4 °C
Spinning

00:05:00 Spin at 500
g

27 Re-suspend pellet in 1 mL ice-cold PBS/BSA 0.04%.

1 mL ice-cold PBS/BSA
0.04%

28 Analyze using hemocytometer with trypan blue. Adjust concentration to 1000 cells / μ L for 10x Chromium or 100 cells / μ L for DropSeq.