VERSION 1

MAY 10, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p2ddqa6

Protocol Citation: Andrew Potter 2018. Adult Mouse Spleen Dissociation (On ice). **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.p2ddqa6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: May 09, 2018

Last Modified: May 10, 2018

PROTOCOL integer ID:
 12069

Keywords: spleen

🌐 Adult Mouse Spleen Dissociation (On ice) V.1

Andrew Potter¹

¹CCHMC

Human Cell Atlas Method Development Community

Andrew Potter

ABSTRACT

Protocol used to dissociate adult (8-10 wk) mouse spleen into single cells. Attained >95% viability, a variety of cell sizes, and ~10 million cells from 12 mg tissue.

GUIDELINES

Collagenase Enzyme Mix (two tubes, 1 mL each)

7.5 mg/mL Collagenase A (Sigma, 10103578001)
 7.5 mg/mL Collagenase Type 4 (Worthington, CLS-4)
 100 µg/mL soybean trypsin inhibitor (Sigma, 10109886001)
 125 U DNase (Applichem, A3778)
 5 mM CaCl₂
 740 µL DPBS (no Ca, Mg)

+12 mg chopped spleen / tube

MATERIALS

MATERIALS

⊗ Red Blood Cell Lysis Buffer Hybri-Max **Sigma Aldrich Catalog #R7757**

BEFORE START INSTRUCTIONS

- Set centrifuges to 4° C.
- Make two tubes of 1 mL enzyme mix.
- Make ~25 mL of DPBS/0.04% BSA

- 1 Chop tissue coarsely (~30 secs) using razor blade on petri dish, on ice.
- 2 Add 12 mg chopped tissue to 1 mL enzyme mix.
- 3 Incubate tube on ice for 10 minutes. Triturate 10X every 2 mins and shake every min.

00:10:00 Incubate on ice
- 4 After 10 mins of digestion, let tissue chunks settle for 1 min on ice & remove 80% of supernatant with released cells & filter using 70 μ M filter on 50 mL conical, on ice. Rinse filter with 5 mL ice-cold PBS/0.04% BSA. Leave filter and 50 mL conical on ice, it will be used for the steps as well.
- 5 Add additional 1 mL enzyme mix to tissue chunks.
- 6 Continue to triturate 10x every 2 minutes and shake every minute while incubating on ice, for 10 additional minutes.

00:10:00 Incubate on ice
- 7 Triturate and add entire volume of cell digestion to 70 μ M filter on 50 mL conical. Rinse w/5 mL ice-cold PBS/0.04% BSA.
- 8 Transfer flow-through to 15 mL conical. Spin 650 g for five minutes at 4 °C. After spin, remove supernatant (down to ~100 μ L).
- 9 Perform RBC lysis: add 1 mL RBC lysis buffer to cells and triturate 10X. Let sit 3 minutes on ice. Add 10 mL ice-cold PBS/BSA 0.04%

00:03:00 incubate on ice

- 10 Spin 650 g for 5 mins at 4 °C. Remove supernatant and re-suspend in 1 mL PBS/BSA 0.04%.