VERSION 2

JUN 18, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.4c3gsyn

Protocol Citation: Andrew Crowley 2019. Agar Plate Preparation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4c3gsyn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 18, 2019

Last Modified: Jun 18, 2019

PROTOCOL integer ID:
 24699

Keywords: bacterial culture, LB, agar

Agar Plate Preparation V.2

Andrew Crowley¹

¹Dartmouth College, Ackerman Lab

Ackerman Lab

Andrew Crowley

Dartmouth College, Ackerman Lab

ABSTRACT

Default scale - 1 L (makes approx. 2.5 sleeves of plates)

MATERIALS

STEP MATERIALS

Agar Alfa Aesar Catalog #A10752-0E

LB Research Products International Corp (RPI) Catalog #L24400-2000.0

Kanamycin Research Products International Corp (RPI) Catalog #K22000-25.0

Ampicillin Research Products International Corp (RPI) Catalog #A40040-25.0

Tryptic Soy Broth Fisher Scientific Catalog #DF0370-07-5

1 Add powdered LB (25 g) and agar (15 g) to deionized water (1 L)

Agar Alfa Aesar Catalog #A10752-0E

LB Research Products International Corp (RPI) Catalog #L24400-2000.0

Note

n.b.

Particulates may remain after addition, but these will dissolve during the autoclave cycle

2 Autoclave using liquid cycle 00:15:00

3 Allow media to cool to 50 °C

4 Select antibiotic from the list below:

Step 4 includes a Step case.

Kanamycin

Ampicillin

step case

Kanamycin

5 Add 50 mg kanamycin and mix thoroughly

- 6 Pour into sterile plates and allow to cool