OCT 24, 2020

Agarose Gel Electrophoresis

Zhujun Wei¹¹Northeast Forest University

2020 iGEM NEFU China

Zhujun Wei

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bnwsmfee

Protocol Citation: Zhujun Wei 2020. Agarose Gel Electrophoresis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bnwsmfee>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 24, 2020

Last Modified: Oct 24, 2020

PROTOCOL integer ID:
43698

ABSTRACT

The concentration of the agarose gel is 1%.

MATERIALS

Agarose
50X TAE
Gelred

SAFETY WARNINGS

Please wear gloves for the experiment, Be careful when heating, protect your hands.

1 Weigh 0.20 g of agarose.

- 2 Add 20 ml of TAE 1X.
Need to prepare 1X TAE with 50X TAE.
- 3 Heat up until the solution is homogeneous.
- 4 Add 1 μ l Gelred into solution and mix well.
- 5 Put the solution into bed for polymerize, make sure "comb" is well placed and the solution is balanced.
- 6 20 min later, use the gel.