JUL 23, 2018

Aiptasia spawning and embryo/larvae handling - Weis Lab

In 1 collection

OSU Weis Lab¹, Jason S Presnell¹

¹Oregon State University

Weis Lab Oregon State

Aiptasia Symbiodiniaceae Model System

Jason S Presnell

Oregon State University

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.rbsd2ne

Protocol Citation: OSU Weis Lab, Jason S Presnell 2018. Aiptasia spawning and embryo/larvae handling - Weis Lab. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.rbsd2ne>

MANUSCRIPT CITATION:

Bucher M, Wolfowicz I, Voss PA, Hambleton EA, Guse A (2016) Development and Symbiosis Establishment in the Cnidarian Endosymbiosis Model Aiptasia sp. *Sci Rep* 6:19867.
Grawunder D, Hambleton EA, Bucher M, Wolfowicz I, Bechtoldt N, Guse A (2015) Induction of Gametogenesis in the Cnidarian Endosymbiosis Model Aiptasia sp. *Sci Rep* 5:15677.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 27, 2018

Last Modified: Jul 23, 2018

PROTOCOL integer ID: 13394

ABSTRACT

This protocol outlines the steps taken to support sexual reproduction in Aiptasia lab cultures. It also provides steps for handling embryos and larvae, and basic husbandry to keep them alive.

Spawning protocol based on the methods from Grawunder et al. 2015.

ATTACHMENTS

[Bucher 2016.pdf](#) [Grawunder 2015.pdf](#)

Pre-spawning

- 1 Aiptasia polyps are kept at low-density cultures to help maximize growth. Typically no more than 3 or 4 polyps are kept in plastic food containers (Cambro 92CW, 17.6 x 10.8 cm). These animals are kept at 25°C under a 12 hour light and dark cycle (standard culture conditions), with white light between 15-20 $\mu\text{mol photons/mm}^2\text{s}$.
- 2 Polyps are fed dense cultures of brine shrimp at least 5 days a week and the water is changed at least every other day. Under this feeding regime, polyps should grow rapidly. Once polyps have grown a sufficient size, i.e. 0.5 cm oral disk diameter, they are placed into a different incubator used for spawning ('spawning incubator'). Polyps continue to be fed brine shrimp at least 5 days a week, with the water changed at least every other day.

- 3 Male and female crosses can be setup in containers, especially if the timing of fertilization is not important and larvae are desired. To control the timing of fertilization, separate sexes are kept in different containers. The light and dark cycles in the spawning incubators can be setup for convenient spawning times. In the lab, Aiptasia typically release gametes 6 hours after the onset of darkness.
- 4 The spawning incubators are setup to mimic the lunar cycle by providing blue light (peak wavelength at 450 nm, intensity $\sim 10 \mu\text{mol photons /mm}^2\text{s}$) every 28 days. The blue light is turned on during the 12 hour dark period for the first 5 days of the 28-day cycle. If well-fed and healthy, polyps will likely spawn within the first two 28-day cycles. Spawning typically occurs between days 13 - 20.

Gamete collection and fertilization

- 5 Eggs sink to the bottom of the dish and are uniformly round, white, and opaque. Eggs are approximately $80 \mu\text{m}$ in diameter and are easily identifiable under a standard stereomicroscope (See image). Sperm is a bit more challenging to observe, but if enough is released the water will appear cloudy.

- 6 Different sizes of mesh screen (e.g. Nitex or cell strainers) can be used to collect gametes. For example, water containing sperm from male-only tanks can be poured through a small mesh screen to filter out large debris. Similarly eggs can be passed through a larger mesh screen (e.g. 100 μm) to filter out large debris.
- 7 The eggs and sperm are then mixed in a 1,500 mL (8-inch) glass culture dish and placed at 25°C. Fertilization should take place within 10-20 minutes, depending on sperm concentration.

Note

Culture dishes are from Carolina Biological Supply Company (https://www.carolina.com/lab-dishes/carolina-culture-dishes/FAM_740996.pr).

Embryo/larvae handling

- 8 Embryos and/or larvae can be concentrated from larger dishes by pouring them through a filter small enough to capture eggs, roughly $< 80 \mu\text{m}$ (e.g., a $40 \mu\text{m}$ cell strainer).
- 9 The captured embryos/larvae can then be placed into a smaller culture dish filled with filtered seawater (FSW). This is a good method to clean the water since ciliates will pass through the filter. Take care to not let the embryos/larvae be exposed to air. Once all the water containing the embryos/larvae has been poured through the filter, immediately immerse the filter in the new culture dish that is filled with FSW. Invert the filter to free the embryos/larvae.
- 10 As noted in (Bucher et al. 2016), early cleavage stages take place a few hours post fertilization (hpf). Embryos reach the blastula stage by 12-14 hpf, and by 24 hpf have undergone gastrulation. Embryos and larvae should be kept at 25°C in 4-inch glass culture dishes (or larger). Changing the water at least once a day should keep the larvae healthy.
- 11 Cilia lining the ectoderm develops and begins to beat by 24 hpf, causing the embryos to "swim" around the bottom of the dish.
- 12 By 48 hpf, planula larvae can be observed actively swimming through the water.