VERSION 1

JUL 04, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.45dgy26

Protocol Citation: Robert Auber, Jennifer H Wisecaver 2019. Algal DNA extraction for HMW Nanopore sequencing. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.45dgy26>

MANUSCRIPT CITATION:

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jul 03, 2019

Last Modified: Jul 04, 2019

PROTOCOL integer ID:
 25477

🌐 Algal DNA extraction for HMW Nanopore sequencing V.1

Robert Auber¹, Jennifer H Wisecaver¹

¹Purdue University

High molecular weight DNA extraction from all kingdoms

Wisecaver Lab

Robert Auber

ABSTRACT

This is a protocol adapted for a haptophyte algae. It is a single cellular organism lacking any type of silica or calcium carbonate armor. If your organism doesn't lyse by a simple snap freeze, some type of grinding step is recommended. Confirmation of properly extracting the nuclei can be done with propidium iodide staining and an epifluorescent microscope. The actual DNA extraction gives ultra long reads (greater than 200kb) by using a circulomics NanoBind kit.

Keywords: Algae, DNA extraction, Nuclei isolation, Nanopore

MATERIALS

Supplies

50 ml conical tubes

LoBind 1.5 ml microcentrifuge tubes

1000 µl and 100 µl wide bore pipette tips

Nylon mesh filters: 100 µm, 70 µm and 40 µm

Liquid nitrogen

Prepared Buffers & Reagents

Nucleus Isolation Buffer (Must be cooled to 0°C for > 1 hour before use):

10 mM Tris pH 9.5

10 mM EDTA 100 mM KCL

500 mM Sucrose

4 mM Spermidine

1 mM Spermine

0.1% BME (add day of extraction)

10% Triton X-100

2-Mercaptoethanol (BME)

Isopropanol

100% Ethanol

Proteinase K (> 600 mAU/mL solution)

RNase A (100mg/mL)

Prepare equipment and reagents

- 1 Set water bath to 74°C
- 2 Chill centrifuge to 4°C
- 3 Add BME to premade NIB (Per Sample 35 µl BME into 35 ml premade NIB + Aliquot 1.8 ml NIB and add 200 µl Triton X-100)

- 4 Cool NIB to 0°C
- 5 Cool NIB + Triton X to 0°C
- 6 Get liquid nitrogen
- 7 Chill four 50 ml conical tubes (per sample) on ice

Cell Lysis

- 8
- 9 Transfer 20-50 ml of culture to a pre-chilled 50 ml conical tube and centrifuge at 4°C at 2000g for 10 min to pellet cells.
- 10 Discard the supernatant and snap freeze the conical tube in liquid nitrogen.
- 11 Place cells on ice and add 10 ml of ice cold NIB.

- 12** Mix by slowly pipetting up and down with a 5 ml pipettor until the mixture is smooth and not clumpy

Filter to remove particulate material

- 13** Place a 100 μm filter on top of a pre-chilled 50 ml conical tube
- 14** Slowly run the sample through the filter into the clean tube
- 15** Wash the filter with 10 ml of ice cold NIB
- 16** Place a 70 μm filter on top of a second pre-chilled 50 ml conical tube
- 17** Slowly run the sample through the filter into the clean tube
- 18** Wash the filter with 10 ml of ice cold NIB
- 19** Place a 40 μm filter on top of a third pre-chilled 50 ml conical tube

- 20 Slowly run the sample through the filter into the clean tube (do not wash with fresh NIB)
- 21 Add 1.5 ml of ice cold NIB+TritonX
- 22 Centrifuge at 4°C at 500g for 10 min to pellet nuclei.
- 23 Discard the supernatant

Extract DNA from Nuclei

- 24 D. Extract DNA by referring to Circulomics Plant Nuclei Big DNA Kit protocol
<https://www.circulomics.com/store/Nanobind-Plant-Nuclei-Big-DNA-Kit-Alpha-Version-p99924200>