FEB 19, 2019

Aliquotting PCR Primer Mixture

Kenneth Schackart¹

¹University of Arizona

Yoon Lab

Kenneth Schackart
University of Arizona

ABSTRACT

How to take lyophilized PCR primers and dilute and aliquot into useful quantities and concentrations.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.yb7fsrn

Protocol Citation: Kenneth Schackart 2019. Aliquotting PCR Primer Mixture.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.yb7fsrn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 19, 2019

Last Modified: Feb 19, 2019

PROTOCOL integer ID:
20575

Dilute primers to 100 μ M

- 1 Add the volume of nuclease-free water to each tube according to the documentation provided by the primer.

In the case of *E. coli* K12, the following volumes are to be used:

Oligo	Volume
F184031	552 μ L
R18551	728 μ L

Make 100 μ L stock

- 2 Use the following table to make 25 μ M, 100 μ L stocks:

Component	Volume
100 μ M F184031	25 μ L
100 μ M R18551	25 μ L
Nuclease Free Water	50 μ L
Total	100 μ L

Making 100 μ L aliquots with 25 μ M of each primer.

- 3 Store all primer mixtures in freezer for up to 2 years

Aliquot primer mixtures

- 4 Aliquot the primer mixture into quantities that can be expected to be used in one day to avoid freeze/thaw cycling.