VERSION 2

APR 27, 2018

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.ptidnke

Protocol Citation: Emil Boros, Marina Kolpakova 2018. An assessment on a large geographic scale of Eurasian inland saline surface waters. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.ptidnke>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Apr 27, 2018

Last Modified: Apr 27, 2018

PROTOCOL integer ID:
11850

An assessment on a large geographic scale of Eurasian inland saline surface waters V.2

Emil Boros¹, Marina Kolpakova²

¹Balaton Limnological Institute, Centre for Ecological Research, Hungarian Academy of Sciences (MTA);

²Sobolev Institute of Geology and Mineralogy, Siberian Branch of Russian Academy of Sciences

Emil Dr Boros

Balaton Limnological Institute, Centre for Ecological Resear...

ABSTRACT

The major ion concentration data of sodium (Na), potassium (K), calcium (Ca), magnesium (Mg), chloride (Cl), sulphate (SO₄), bicarbonate (HCO₃), carbonate (CO₃) and pH (if it was coupled with ion data) were input into the database. The data were drawn from a large geographic scale across Eurasia (Austria, China, Hungary, Kazakhstan, Mongolia, Russia, Serbia, Turkey) and from a large number of saline lakes and pans (N=220) with minimum a 1.0 g L⁻¹ salinity threshold. The 1.0 g L⁻¹ salinity threshold was selected based on a former study (Boros et al., 2014), where this threshold was experimentally found to be the characteristic boundary of soda ecosystems. Salinity was estimated by the sum of measured concentrations of eight major ions (Na, K, Ca, Mg, Cl, SO₄, HCO₃, CO₃). As generally known, sodium is by far the most common cation in saline lakes and necessarily the dominant cation in soda type lakes. Therefore, sites were excluded from the dataset if Na was not the most abundant ion. If seasonal or annual water data were available, mean values were put into database. Most of the data came from papers (sources are indicated in the table).

Keywords: alkaline,
bicarbonate, carbonate,
chemical types of saline
waters, equivalent
percentage (e%), pH overlap,
rank of dominant ion, Saline
type, Soda type, Soda-Saline
type, soda brine

1

repository_data_ebmk_final.xlsx