



Aug 15, 2023

Anatomical variations and dimensions of the popliteus muscle in cadaveric specimens

 [F1000Research](#)

DOI

dx.doi.org/10.17504/protocols.io.3byl4qqk8vo5/v1

Murlimanju Virupakshamurthy¹, Rajanigandha Vadgaonkar¹

¹Department of Anatomy, Kasturba Medical College, Mangalore, Manipal Academy of Higher Education, Manipal, India



Murlimanju Virupakshamurthy

Department of Anatomy, Kasturba Medical College, Mangalore, ...

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.3byl4qqk8vo5/v1>

Protocol Citation: Murlimanju Virupakshamurthy, Rajanigandha Vadgaonkar 2023. Anatomical variations and dimensions of the popliteus muscle in cadaveric specimens. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3byl4qqk8vo5/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 15, 2023

Last Modified: August 15, 2023

Protocol Integer ID: 86489

Keywords: morphology of the popliteus muscle, morphology of popliteus muscle, popliteus muscle, biomechanics of popliteus, dimensions of the popliteus muscle, popliteus muscle spasticity, lateral collateral ligament, lateral femoral condyle, knee joint, fibular ligament, ankle joint, insertion of popliteus, involvement of popliteus, tendon, actions on the knee joint, origin of popliteus, morphometric data of popliteus, femur bone, styloid process of the fibula, rotation of the tibia, complex injury of the knee joint, lateral meniscus, popliteus, knee, popliteus at various location, tibial nerve, adult human cadaveric lower limb, tear of lateral meniscus, tendon in context, femur, sided lower limb, lower limb, fibula, anatomical variation, additional stress on the tendon, unlocking muscle, popliteo, leg, biomechanics, muscle belly, back of leg, anatomy, only muscle, posterolateral part, visible deformity, tibia, intact posterior compartment, morphology

Abstract

Introduction

The popliteus muscle is located at the flexor aspect of the leg and is supplied by the tibial nerve. This is the only muscle in the back of leg, which actions on the knee joint and never over the ankle joint. This is considered as the knee's unlocking muscle. It laterally swaps the femur over tibia, while walking when one foot is on the ground. It also causes medial rotation of the tibia on the femur bone, once the foot is over the ground. Along with lateral collateral ligament and popliteo-fibular ligament, the tendon of popliteus plays instrumental role in the stabilization of the posterolateral part of knee (1). The lateral femoral condyle and lateral meniscus at its posterior horn, offers the origin of popliteus. Its origin is tendinous and it is interesting to know that there exists variability in its origin like from the styloid process of the fibula (2).

Aims and Objective

The goal of this investigation is to describe the morphology of popliteus muscle with its tendon in context to its variability in origin, mode of insertion, innervation patterns and its dimensions. The objectives are to measure the length, thickness and width of the popliteus at various locations.

Methodology

Study setting: Department of Anatomy, Kasturba Medical College, Mangalore

Study design: Institutional Based Descriptive Cross Sectional Study

Study participants: A meticulous dissection will be done in adult human cadaveric lower limb specimens and the morphology of the popliteus muscle will be tabulated for the statistical analysis.

Inclusion criteria: Lower limb specimens with an intact posterior compartment will be included

Exclusion criteria: The specimens showing fractures, any visible deformities and missing parts will be excluded.

Study duration: 3 months

Sample size: 25 adult cadavers (25 right and 25 left sided lower limbs)

Sampling method: The sample size is similar to the earlier study performed by Olewnik et al. (1).

Data collection methodology: The variability of the origin of popliteus and the insertion of popliteus will be noted down. To study the morphometry, the popliteus muscle is divided into certain sections extending from the origin to its final insertion into the posterior surface of the tibia. The parameters considered in the study are measuring the length, width and thickness of the muscle belly and its tendon, which will be taken by digital Vernier caliper.

Data analysis: The data collected as above will be analyzed by applying paired samples t test.



Implications

The anatomy and biomechanics of popliteus makes it an important structure, which keeps the knee stable. But its involvement is ignored in the complex injury of the knee joint (3). The isolated involvement of popliteus is seen in sports injuries and it may be misinterpreted as a tear of lateral meniscus. The sports like tennis, basketball and downhill running may put additional stress on the tendon of popliteus (3). In this context, the morphometric data of popliteus are clinically important for the effective treatment of the popliteus muscle spasticity (4).

Troubleshooting

Protocol references

1. Olewnik, Ł., LaPrade, R.F., Paulsen, F. *et al.* A proposal for a new morphological classification of the popliteus muscle tendon with potential clinical and biomechanical significance. *Sci Rep* **11**, 14434 (2021).
1. Hyland S, Varacallo M. **Anatomy, Bony Pelvis and Lower Limb, Popliteus Muscle.** [Updated 2020 Aug 15]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020.
1. Zabrzyński J, Huri G, Yataganbaba A, Paczesny Ł, Szwedowski D, Zabrzyńska A, Łapaj Ł, Gagat M, Wiśniewski M, Pękala P. Current concepts on the morphology of popliteus tendon and its clinical implications. *Folia Morphol (Warsz)*. 2021;80(3):505-513.
1. Jeon A, Kim YG, Sohn Y, Lee JH. Nerve and Arterial Supply Pattern of the Popliteus Muscle and Clinical Implications. *Biomed Res Int*. 2022 Jan 10;2022:6980471.