

Aug 10, 2022

Anthoceros punctatus (hornwort) transformation

 In 1 collection

DOI

dx.doi.org/10.17504/protocols.io.81wgb6jz3lpk/v1

Eftychis Frangedakis¹, Manuel Waller²


¹University of Cambridge; ²University of Zürich

Hornworts



Eftychis Frangedakis

University of Cambridge, Plant Sciences



Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.81wgb6jz3lpk/v1>

Protocol Citation: Eftychis Frangedakis, Manuel Waller 2022. Anthoceros punctatus (hornwort) transformation. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.81wgb6jz3lpk/v1>

Manuscript citation:

<https://doi.org/10.1101/2022.08.10.503456>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 03, 2022

Last Modified: August 10, 2022

Protocol Integer ID: 68157

Keywords: transformation anthoceros punctatus, anthoceros punctatus, hornwort, transformation

Abstract

Anthoceros punctatus (hornwort) transformation



Materials

KNOP recipe:

Stock 1

25g/L KH_2PO_4

Stock 2

25g/L KCl

Stock 3

25g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$

Stock 4

100g/L $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$

autoclave and store at RT or 4°C

KNOP solid working solution:

In 600 mL of water add:

10ml Stock 1

10ml Stock 2

10ml Stock 3

10ml Stock 4

12.5mg $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

pH to 5.8 with KOH

top up water to 1L after adjusting pH

add 7 gr of Gelzan - G1910 - CAS Number [71010-52-1](#)

KNOP liquid working solution:

In 600 mL of water add:

10ml Stock 1

10ml Stock 2

10ml Stock 3

10ml Stock 4

12.5mg $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$

10 gr of sucrose (1% w/v final concentration)

40mM MES (*very important*)

pH to 5.5 with KOH

top up water to 1L after adjusting pH

Filter sterile (do not autoclave), aliquot into 50mL falcon tubes and store at -20°C.

Sterile disposable scalpels (#0501, Swann Morton)

Razor blades (#11904325, Fisher Scientific)

100 µm cell strainer (#352360, CORNING),

6-well plate (#140675, ThermoFisher)

3',5'-dimethoxy-4'-hydroxyacetophenone (acetosyringone) (#115540050, Acros Organics, dissolved in dimethyl sulfoxide (DMSO) (#D8418, SIGMA))

Cefotaxime (#BIC0111, Apollo Scientific)

Hygromycin (#10687010, Invitrogen)

Corning Disposable Vacuum Filter/Storage Systems (#430767)

Troubleshooting

1 Axenic cultures of *A. punctatus* gametophytes can be routinely propagated by monthly sub-culturing

2



Figure 1: 6 week old *A. punctatus* thallus

3




Figure 2: Transformation method outline. A) Thallus tissue is routinely propagated on a monthly basis under low light. 4-6 week old tissue is fragmented with the aid of a razor blade, transferred to a cell strainer, and washed thoroughly with sterile water. B) The tissue is then co-cultivated with *Agrobacterium* for three days (under low light) and C) spread on antibiotic-containing growth medium. After approximately 4-6 weeks, putative transformants are visible. A final round of selection is recommended to eliminate false-positive transformants.

4 Tissue preparation:

- Collect approximately 1 g of thallus tissue grown for 4-6 weeks under low light intensity (approximately 0.1 g of tissue per petri dish - 10 petri dishes in total). **Figure1 and Figure 3.1**
- Transfer the tissue into an empty petri dish, add sterile water until the tissue is covered and fragment using a razor blade (it takes approximately 5 mins, similar to Video 1). **Figure 3.2-3**

- Transfer the tissue from the petri dish into a cell strainer positioned on a falcon tube using sterile scalpels and wash the tissue using ~100 ml of sterile water or until the flow through was clear. **Figure 3.4-6**

5 **Agrobacterium culture preparation:**

- Inoculate 5 mL LB media with 3-4 *Agrobacterium* colonies (*AGL1*: 15 µg/mL rifampicin, 50 µg/mL carbenicillin) (*GV3101*: 50 µg/mL rifampicin, 25 µg/mL gentamicin) and the plasmid-specific selection antibiotic.
- Incubate the preculture at 28°C for 2 days at 110 rpm.
- Centrifuge 5 mL of 2 d *Agrobacterium* culture (no need to measure OD) for 7 min at 2000 xg.
- Remove supernatant and re-suspend in 5 mL liquid KNOP plus 1% (w/v) sucrose and 100 µM acetosyringone.
- Incubate the culture with shaking (120 rpm) at 28°C for 3-5 hours.

6 **Co-cultivation:**

- Transferred the fragmented thallus tissue into a 6-well plate (transfer 1/6 of the 1 g tissue into a single well) with 5 mL of liquid KNOP medium supplemented with 1% (w/v) sucrose and **30-40 mM MES (VERY IMPORTANT), pH 5.5**, 80 µL of *Agrobacterium* culture and acetosyringone at final concentration of 100 µM. **Figure 3.7**
- Co-cultivate the tissue with the *Agrobacterium* for 3 days on a shaker at 110 rpm, with only ambient light.
- Using a sterile plastic pipette transfer the tissue of one well into a cell strainer, drain and then transfer on growth media containing the appropriate antibiotic (onto 1 petri dish from one well). To facilitate spreading of the tissue, 1-2 mL of sterile water is added to the petri dish. **Figure 3.8-11**
- After 6-7 weeks successful transformants are visible on the petri dish (successful transformants can be identified using a dissecting scope after 4 weeks selection (sometimes as early as 2 weeks) based on rhizoid production and/or fluorescence if such a marker is present on the construct). **Figure 3.12**

- The emergence of rhizoids is an indication of successful transformation (yellow arrow: transformed thallus fragment, blue arrow: dying thallus fragment). **Figure 4**

7

2nd selection (necessary):

- To eliminate false positives, after 4 weeks transfer the tissue to fresh growth media containing 100 µg/mL cefotaxime and 10 µg/mL Hygromycin. To facilitate spreading of tissue on the petri dish add 2 mL of sterile water. Grow at 21°C under 12 hours light and 12 hours dark, 35 µmol m⁻²s⁻¹. Sometimes a 3rd round of selection is necessary.

8





Figure 3

9



Figure 4: The emergence of rhizoids is an indication of successful transformation (yellow arrows: transformed thallus fragment, blue arrow: dying thallus fragment).

10



Example of plate with successful transformants 8 weeks after co-cultivation.

11





Video 1, Example of tissue fragmentation for *A. agrestis* Bonn. The method is the same for *A. punctatus*.