Anti-BrdU Staining Using 70% Ethanol and 2N HCl V.3

Kelsey Miller¹

¹BioLegend

Sam Li
BioLegend

ABSTRACT

Anti-BrdU Staining Protocol Using 70% Ethanol and 2N HCl

DOI:

dx.doi.org/10.17504/protocols.io.tibekan

External link:

http://www.biolegend.com/media_assets/support_protocol/BrdU_Staining_Protocols_05102016.pdf

Protocol Citation: Kelsey Miller . Anti-BrdU Staining Using 70% Ethanol and 2N HCl. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.tibekan>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Sep 12, 2018

Last Modified: Aug 29, 2019

PROTOCOL integer ID:
15651

- 1 Pulse actively dividing cells with BrdU (in vitro, cell culture media can be pulsed by adding 10-40 μ M of BrdU for 1-2 hours).
- 2 Harvest cells and centrifuge for 5 minutes at 1200-1500 rpm (200-300 x g)

00:05:00

- 3 Wash cells in 1x PBS (PBS, 10x Concentrate, Cat. No. 926201) and centrifuge for 5 minutes at 1200-1500 rpm (200-300 x g). Discard supernatant.

NOTE: The combined presence of proteins and HCl in downstream steps may cause aggregation. As such, it is highly recommended that wash steps utilize PBS without any protein additive until otherwise indicated

00:05:00

- 4 Dislodge cell pellet and add 5 ml of ice-cold (-20°C) 70% Ethanol to $1-2 \times 10^7$ cells dropwise while slowly vortexing. Incubate at -20°C for at least 2 hours. Cells may be stored for several days.

Note: For ethanol permeabilization, we recommend this step last for at least 2 hours as this improves permeabilization and overall staining. If you're still having difficulty, you can attempt to extend this to an overnight incubation (assuming you aren't also staining for cell surface markers that could be lost).

02:00:00

- 5 Repeat step 3 twice

- 6 Dislodge cell pellet and add 2 ml of 2 N HCl and incubate for 20 minutes at room temperature

00:20:00

- 7 Repeat step 3.

- 8 Dislodge cell pellet and add 2ml of 0.1M $\text{Na}_2\text{B}_4\text{O}_7$ for 10 minutes at room temperature.

00:10:00

- 9 Repeat step 3.

- 10 Resuspend cells at a concentration of 1×10^7 cells per/ml of staining buffer and aliquot 100µl per tube. Add anti-BrdU antibody at appropriate concentration and incubate for 20 minutes at room temperature.

00:20:00

- 11** Wash cells in Cell Staining Buffer (Cat. No. [420201](#)) and centrifuge for 5 minutes at 1200-1500 rpm (200-300xg).

00:05:00

- 12** Stain DNA by adding 1µg of either 7-AAD (Cat. No. [420403](#)) or DAPI (Cat. No. [422801](#)). Wait for 5 minutes prior to acquiring samples on flow cytometer.

Note: Adding a 7-AAD or DAPI stain allows you to analyze total DNA content and provides the characteristic horseshoe flow cytometric staining pattern when compared against BrdU. This helps identify the different phases of the cell cycle.