



May 17, 2020

Antibiotic concentrations for LB agar plates

DOI

dx.doi.org/10.17504/protocols.io.bghkjt4w

Yoan Coudert¹

¹Ecole Normale Supérieure de Lyon

CoudertLab



Yoan Coudert

Ecole Normale Supérieure de Lyon

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bghkjt4w

Protocol Citation: Yoan Coudert 2020. Antibiotic concentrations for LB agar plates. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.bghkjt4w>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 17, 2020

Last Modified: May 17, 2020

Protocol Integer ID: 37132



1

| Commonly Used Antibiotics | Recommended Concentration |
|---------------------------|---------------------------|
| Ampicillin | 100 µg/mL |
| Bleocin | 5 µg/mL |
| Carbenicillin | 100 µg/mL |
| Chloramphenicol | 25 µg/mL |
| Coumermycin | 25 µg/mL |
| Gentamycin | 10 µg/mL |
| Kanamycin | 50 µg/mL |
| Spectinomycin | 50 µg/mL |
| Tetracycline | 10 µg/mL |

Stock solutions = 1000X (e.g. Amp stock = 1g in 10ml milliQ water)