VERSION 1

JUN 13, 2018

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.p9adr2e

Protocol Citation: Diep R Ganguly, Peter Crisp 2018. Arabidopsis leaf protoplasting. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p9adr2e>

MANUSCRIPT CITATION:

Crisp P.A., Ganguly D.R., Smith A.B., Murray K.D., Estavillo G.M., Searle I., ... Pogson B.J. (2017) Rapid Recovery Gene Downregulation during Excess-Light Stress and Recovery in Arabidopsis. The Plant Cell 29, 1836–1863.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We have used this protocol to successfully measure gene expression in protoplasts under stress conditions.

🌐 Arabidopsis leaf protoplasting V.1

Diep R Ganguly¹, Peter Crisp²

¹University of Pennsylvania; ²Australian National University

Pogson Group

Diep R Ganguly

University of Pennsylvania, The Australian National University

ABSTRACT

Method for isolating mesophyll protoplasts from Arabidopsis as a system for monitoring RNA stability using transcriptional inhibitors (e.g. cordycepin) under stress treatments (e.g. high-light or hydrogen peroxide treatment).

Adapted from Yoo S.-D., Cho Y.-H. & Sheen J. (2007) Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. Nature Protocols 2, 1565–1572.

GUIDELINES

Adapted from Yoo S.-D., Cho Y.-H. & Sheen J. (2007) Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. Nature Protocols 2, 1565–1572.

Ideally, preps will be done on 3-4 week old *Arabidopsis* plants (before flowering). Choose well-expanded leaves (usually #5 & #7).

Protoplasting 80 leaves (using ~ 65 - 75 mL enzyme solution) from 3 week old plants yielded approx. 1.5 million protoplasts (diluted to 20,000/mL) from which 1.4 - 2.8 ug total RNA (of high quality) could be extracted using TRIzol.

Higher RNA yields were obtained using fewer leaves and more enzyme solution (40-50 leaves of 3 week old plants gave ~4 ug total RNA).

Created: May 20, 2018

Last Modified: Jun 13, 2018

PROTOCOL integer ID:
12290

Keywords: protoplasting

MATERIALS

MATERIALS

- ☒ D-Mannitol **Sigma Aldrich**
- ☒ Vacuum system **Contributed by users**
- ☒ KCl **Contributed by users**
- ☒ CaCl₂ **Contributed by users**
- ☒ MgCl₂ **Contributed by users**
- ☒ NaCl **Sigma Aldrich Catalog #53014**
- ☒ BSA **Sigma Aldrich Catalog #A7906**
- ☒ b-mercaptoethanol **Sigma-aldrich Catalog #M3148-25ML**
- ☒ MES, sodium salt **Bio Basic Inc. Catalog #MB0611.SIZE.250g**
- ☒ Syringe Filtration - Aqueous Solutions, Sterile, Individual Pack, 0.45UM, 13mm, 10/Pk **Bio Basic Inc. Catalog #SFA1345S.SIZE.1**
- ☒ Macerozyme R-10 **Gold Biotechnology Catalog #M8002**
- ☒ Razor blades **Fisher Scientific Catalog #12-640**
- ☒ Centrifuge **Contributed by users**
- ☒ Compound Microscope **Contributed by users**
- ☒ Cellulase R10 **Contributed by users**
- ☒ BRAND® counting chamber BLAUBRAND® Neubauer improved New without clip double ruled **Sigma Catalog #BR717805**
- ☒ Round bottom glass centrifuge tubes **Contributed by users**
- ☒ Miracloth **Merck Millipore Catalog #475855**

BEFORE START INSTRUCTIONS

Before starting experiment, it is considered ideal to keep intended *Arabidopsis* plants for protoplast extraction in the dark overnight. This is so that all the starch is used up, making the isolation procedure easier (less junk in extract).

Prepare the following solutions:

Enzyme Solution: Make solution of 20mM MES (pH 5.7), 0.4M Mannitol and 20mM KCl. Heat solution to 70°C for 3-5min. To this add cellulose R10 (1.5%) and macerozyme R10 (0.4%). Warm solution to 55°C for 10 min (inactivates DNase and proteases and aids solubilising enzyme). Cool to room temperature and add 10mM Ca Cl₂, 1-5mM β-mercaptoethanol (optional) and 0.1% BSA. Filter the final enzyme solution through a 0.45um syringe filter into a petri dish (enzyme solution should be made fresh, however, can be stored overnight at 4°C).

W5 Solution: 2mM MES (pH 5.7), 154mM NaCl, 125mM CaCl₂ and 5mM KCl.

MMG Solution: 4mM MES (pH5.7), 0.4M Mannitol and 15mM MgCl₂.

Keep plants in dark overnight (the day before)

- 1 Before starting experiment, it is considered ideal to keep intended *Arabidopsis* plants for protoplast extraction in the dark overnight. This is so that all the starch is used up, making the isolation procedure easier (less junk in extract).

Use 3-4 week old *Arabidopsis* plants (before flowering). Choose well-expanded leave (usually #5 & #7). Approx. 40 leaves (in 50ml enzyme solution) should give a yield of roughly 1, 500, 000 protoplasts (can extract almost 3ug RNA from this using TRIzol method). This is suitable for routine experiments. For larger scale: Sheen et al. suggests 100 – 150 leaves in 40 – 60ml enzyme solution yielding 10^7 .

Prepare buffers (the day before)

- 2 **Enzyme Solution:** Make solution of 20mM MES (pH 5.7), 0.4M Mannitol and 20mM KCl. Heat solution to 70°C for 3-5min. To this add cellulose R10 (1.5%) and macerozyme R10 (0.4%). Warm solution to 55°C for 10 min (inactivates DNase and proteases and aids solubilising enzyme). Cool to room temperature and add 10mM CaCl_2 , 1-5mM β -mercaptoethanol (optional) and 0.1% BSA. Filter the final enzyme solution through a 0.45um syringe filter into a petri dish (enzyme solution should be made fresh, however, can be stored overnight at 4°C).

W5 Solution: 2mM MES (pH 5.7), 154mM NaCl, 125mM CaCl_2 and 5mM KCl.

MMG Solution: 4mM MES (pH5.7), 0.4M Mannitol and 15mM MgCl_2 .

Enzyme digestion of leaf material

- 3 Cut 0.5-1mm leaf strips of the leaf using a fresh sharp razor blade (on a nice cutting surface) without tissue crushing at the cutting site (i.e. try not to push down on the tissue with the blade, rather try to “glide” the blade to make the cut).
- 4 Transfer leaf cuts quickly and gently into enzyme solution and make sure leaf strips are submerged nicely.
- 5 Vacuum infiltrate 3 x 10 min.

- 6 Continue digestion, **without shaking**, in the dark for at least 3 hours @ room temperature (enzyme solution should turn from a light brown colour to green after gentle swirling which indicates protoplast release).
- 7 Check for release of protoplasts in solution under the microscope (after gentle swirling). *Arabidopsis* mesophyll protoplasts are approximately 30 - 50 µm in diameter.

Resuspend protoplasts

- 8 Dilute the enzyme/protoplast solution with an equal volume of **W5 solution**.
- 9 Rinse 2 clean layers of miracloth in **W5**.

Filter the enzyme solution containing the protoplasts with these layers of miracloth.
- 10 Centrifuge the flow-through at 150 g/rcf for 2 minutes (brake ON/hard stop) in a 30ml-round-bottom glass centrifuge tube.

Remove as much supernatant as possible and re-suspend protoplasts by gentle swirling.
- 11 Use haemocytometer to calculate the concentration of protoplasts.
- 12 Re-suspend in **W5 solution** to a concentration of 2×10^5 /mL.
- 13 Rest protoplasts on ice for at least 30 minutes. Protoplasts should begin to settle to the bottom of the tube by gravity. Remove as much W5 solution as possible without touching the pellet. Re-suspend at 2×10^5 /mL in **MMG solution** kept at room temperature (will need to do another count using haemocytometer before resuspension).