VERSION 1

FEB 10, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.brdfm23n

Protocol Citation: IIDP-HIPP 2021. Assessment of Human Islet Composition and Acinar Cell Component by Immunofluorescence Staining. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.brdfm23n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jan 12, 2021

Last Modified: Feb 10, 2021

PROTOCOL integer ID:
 46215

Assessment of Human Islet Composition and Acinar Cell Component by Immunofluorescence Staining V.1

IIDP-HIPP¹

¹Integrated Islet Distribution Program and Human Islet Phenotyping Program

Integrated Islet Distribution Program and Human Islet Phenotyping Program

Tech. support email: heather.durai@vumc.org

IIDP-HIPP

ABSTRACT

This Standard Operating Procedure (SOP) is based on the Vanderbilt Human Islet Phenotyping Program Core Facility (VHIPP) Immunofluorescence Staining Procedure. This SOP provides HIPP procedure for immunofluorescent staining, imaging, and analysis of islet preparations.

This SOP defines the assay method used by the Human Islet Phenotyping Program (HIPP) for quantitative and qualitative determination of the Purified Human Pancreatic Islet product, post-shipment, manufactured for use in the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK)-sponsored research in the Integrated Islet Distribution Program (IIDP).

Note

This Standard Operating Procedure (SOP) #: HIPP-07-v02

GUIDELINES

- ***Integrated Islet Distribution Program (IIDP) (RRID:SCR_014387):*** The IIDP is a grant funded program commissioned and funded by the NIDDK to provide quality human islets to the diabetes research community to advance scientific discoveries and translational medicine. The IIDP consists of the NIDDK Project

Keywords: Human Islet Composition, Acinar Cell Component, Immunofluorescence Staining, HIPP, IIDP

Scientist and Program Official, the External Evaluation Committee and the CC at City of Hope (COH). The IIDP CC integrates an interactive group of academic laboratories including the subcontracted IIDP centers.

- ***IIDP Coordinating Center (CC):*** Joyce Niland, Ph.D. and Carmella Evans-Molina, M.D., Ph.D. serve as Co-Principal Investigators (Co-PIs) for the IIDP Program located within the Department of Diabetes and Cancer Discovery Science at COH to coordinate the activities of the IIDP and Human Islet Phenotyping Program (HIPP). Dr. Niland, contact PI, oversees the daily activity of the IIDP staff, provides informatics/ biostatistical input, and subcontracts with the Islet Isolation Centers (IICs) to ensure the delivery of the highest quality human islets to IIDP-approved investigators. Dr. Evans-Molina serves as the liaison to the HIPP, interacting closely to ensure that extensive, high quality phenotypic data are collected on islets distributed by the IICs. She also facilitates the delivery of this information to both the IICs and the IIDP-approved investigators, while responding to questions, issues, or suggestions for further HIPP enhancements.
- ***Human Islet Phenotyping Program (HIPP):*** The HIPP is a subcontracted entity of the IIDP through the COH and Vanderbilt University. The HIPP is directed by Marcela Brissova, Ph.D. and is responsible for performing specific standardized phenotyping assays agreed upon by both the IIDP and the HIPP, in order to provide enhanced, quality data on the human islets post-shipment, to the IIDP. The results of these assays will be approved by the CC and posted on the IIDP website for both the centers and the approved investigators.
- ***Cryosections:*** Sections of a tissue/cells embedded in optimal cutting temperature (OCT) compound and frozen -80 °C .
- ***Indirect Immunofluorescence Staining:*** Immunohistochemical procedure based on antigen detection by fluorescence in histological sections using a combination of primary and secondary antibodies where the primary antibody is directed to the antigen of interest and the fluorescently-conjugated secondary recognizes species where the primary antibody was raised. Histological sections are viewed using a microscope system equipped with an appropriate light source and filter set to allow for visualization of fluorescence tissue staining.

References:

CITATION

Dai C, Brissova M, Hang Y, Thompson C, Poffenberger G, Shostak A, et al. Islet-enriched gene expression and glucose-induced insulin secretion in human and mouse islets. *Diabetologia*. 2012 Mar;55(3):707–18. PMCID: PMC3268985.

LINK

<https://pubmed.ncbi.nlm.nih.gov/22167125/>

CITATION

Guo S, Dai C, Guo M, Taylor B, Harmon JS, Sander M, et al. Inactivation of specific β cell transcription factors in type 2 diabetes. *J Clin Invest*. 2013 Aug;123(8):3305–16. PMCID: PMC3726150.

LINK

<https://pubmed.ncbi.nlm.nih.gov/23863625/>

MATERIALS

1. PBS (phosphate buffered saline) with no Ca/Mg, 1X (Invitrogen 14190-144)
2. BSA (bovine serum albumin, Sigma A-6003)
3. SlowFade Gold (Molecular Probes S36938)
4. Triton X-100 (BioRad 1610407)
5. Normal Donkey Serum (NDS, Jackson Immuno Research 017-000-121)
6. 4',6-diamidino-2-phenylindole (DAPI, ThermoFisher Scientific D1306)
7. Kartell Staining Chambers (VWR 25460-907)
8. PAP Marker (Research Products International 195506)
9. 4 °C Refrigerator (ArcticTemp)

PBS (phosphate buffered saline) with no Ca/Mg 1X **Thermo Fisher Scientific Catalog #Invitrogen 14190-144**

BSA (bovine serum albumin) **Merck MilliporeSigma (Sigma-Aldrich) Catalog #A-6003**

SlowFade Gold (Molecular Probes) **Thermo Fisher Scientific Catalog #S36938**

Triton X-100 **Bio-Rad Laboratories Catalog #1610407**

Normal Donkey Serum **Jackson ImmunoResearch Laboratories, Inc. Catalog #017-000-121**

Equipment

Kartell Staining Chambers

NAME

VWR

BRAND

25460-907

SKU

<https://us.vwr.com/store/product/4619254/vwr-staining-jar-with-cover-hellendhal-type>

LINK

Equipment

Marker

NAME

PAP

BRAND

195506

SKU

<https://www.rpicorp.com/products/molecular-biology/hybridization/super-ht-mini-pap-pen-2-5mm.html>

LINK

Equipment

4°C Refrigerator

NAME

ArcticTemp

BRAND

None

SKU

https://www.meatprocessingproducts.com/arctic-air-commercial-refrigerators.html?gclid=EAlaIQobChMI-lzMrrKo7gIVAoIGCh2-zAxFEAAYASAAEgJWDvD_BwE

LINK

SAFETY WARNINGS

Triton X-100 (*BioRad 1610407*)

Safety information

Triton X-100.pdf

Preparation of Reagents

1 Preparation of Reagents for Immunofluorescence Staining of Islet Cryosections

- 1.1 10% Triton X-100 stock** (30 mL) – combine 3 mL Triton-X-100 and 27 mL 1X PBS. Mix on shaker for 30 min or until Triton X-100 is completely dissolved and store at 4 °C for up to 1 month.

- 1.2 **Permeabilization Solution (0.2% Triton, 50 mL)** – combine 1 mL of 10% Triton stock and 49 mL 1X PBS.
- 1.3 **Blocking Buffer (5% NDS, 4 mL)** – combine 0.2 mL NDS and 3.8 mL 1X PBS.
- 1.4 **Antibody Buffer (10 mL)** – combine 0.1 g BSA, 0.1 mL 10% Triton stock, and 9.8 mL 1X PBS.
- 1.5 **DAPI staining solution (1:25,000, 50 mL)** – combine 2 µL DAPI stock (5mg/mL) and 50 mL 1X PBS.

Procedure

2 Immunofluorescence Staining Procedure on Islet Cryosections

- 2.1 Use freshly-made antibody incubation buffers, and wash buffers. Steps 2.3, 2.4, 2.8, 2.10, 2.11 can be done in Kartell Staining Chambers.
- 2.2 Let the frozen sections thaw at room temperature and air-dry for about 30 minutes.
- 2.3 Wash the sections with 50 mL 1X PBS 3 times for 5 minutes to remove the OCT.

- 2.4** Permeabilize the tissue section with 0.2 % Triton for 15 minutes at room temperature.
- 2.5** Wash the tissue in 50 mL 1X PBS 3 times for 3-5 minutes.
- 2.6** Draw circles or rectangles around the sections with PAP marker and let them dry for about 5 minutes.
- 2.7** Block the sections with 5% normal donkey serum (made from 100% stock)/1X PBS at room temperature for 90 minutes in a humidified chamber.
- 2.8** Aspirate the blocking solution, add primary antibodies (**Table 1**) diluted in 0.1% Triton-X-100 (made from 10% Triton stock)/1% BSA/1X PBS and incubate in a humidified chamber overnight at 4 °C .

Table 1. List of primary and secondary antibodies for assessment of islet cell composition and endocrine/acinar cell composition

Primary Antibody	Vendor	Product number	Final Dilution	Secondary Antibody	Vendor	Product number	Final Dilution
C-peptide (rat)	Developmental Studies Hybridoma Bank	GN-ID4 RRID:AB_2631151	1:100	Rat IgG-Cy2 (donkey)	Jackson ImmunoResearch	712-225-150 RRID:AB_2340673	1:500
				Rat IgG-Cy5 (donkey)	Jackson ImmunoResearch	712-175-150 RRID:AB_2340671	1:200
Glucagon (mouse)	Abcam	ab10988 RRID:AB_297642	1:250	Mouse IgG-Cy3 (donkey)	Jackson ImmunoResearch	715-165-150 RRID:AB_2340813	1:500
Glucagon (rabbit)	Cell Signaling Technology	2760S RRID:AB_659831	1:100	Rabbit IgG-Cy5 (donkey)	Jackson ImmunoResearch	711-175-152 RRID:AB_2340607	1:200
Somatostatin (goat)	Santa Cruz Biotechnology	sc-7819 RRID:AB_2302603	1:500	Goat IgG-Cy5 (donkey)	Jackson ImmunoResearch	705-175-147 RRID:AB_2340415	1:200
HPX1 (mouse)	Novus Biologicals	NBP1-18951 RRID:AB_1625456	1:100	Mouse IgG-Cy3 (donkey)	Jackson ImmunoResearch	715-165-150 RRID:AB_2340813	1:500

Note

Antibody Information

Primary Antibodies:

C-peptide (rat): GN-ID4 [RRID:AB_2631151](https://rrid.org/RRID:AB_2631151)

Glucagon (mouse): ab10988 [RRID:AB_297642](https://rrid.org/RRID:AB_297642)

Glucagon (rabbit): 2760S [RRID:AB_659831](https://rrid.org/RRID:AB_659831)

Somatostatin (goat): sc-7819 [RRID:AB_2302603](https://rrid.org/RRID:AB_2302603)

HPX1 (mouse): NBP1-18951, [RRID:AB_1625456](https://rrid.org/RRID:AB_1625456)

Secondary Antibodies:

Rat IgG-Cy2 (donkey): 712-225-150 [RRID:AB_2340673](https://rrid.org/RRID:AB_2340673)

Rat IgG-Cy5 (donkey): 712-175-150 [RRID:AB_2340671](https://rrid.org/RRID:AB_2340671)

Mouse IgG-Cy3 (donkey): 715-165-150 [RRID:AB_2340813](https://rrid.org/RRID:AB_2340813)

Rabbit IgG-Cy5 (donkey): 711-175-152 [RRID:AB_2340607](https://rrid.org/RRID:AB_2340607)

Goat IgG-Cy5 (donkey): 705-175-147 [RRID:AB_2340415](https://rrid.org/RRID:AB_2340415)

Mouse IgG-Cy3 (donkey): 715-165-150 [RRID:AB_2340813](https://rrid.org/RRID:AB_2340813)

- 2.9 Aspirate the primary antibodies and wash the sections with 1X PBS three times for 10 minutes each.
- 2.10 Add secondary antibodies (**Table 1**) diluted in 0.1% Triton/1% BSA/1X PBS and incubate for 1.5 hours at room temperature in a humidified chamber.
- 2.11 Aspirate the secondary antibody and counterstain slides with 1:25,000 DAPI/PBS for 10 minutes at room temperature.
- 2.12 Remove from DAPI and wash the sections with 1X PBS three times for 15 minutes each.
- 2.13 Mount the sections with SlowFade Gold mounting medium.

Imaging and Analysis

3 Imaging and Analysis of Fluorescently Labeled Islet Cryosections

- 3.1 Capture images of islet sections using a high-resolution whole slide scanning system (ScanScope FL, Aperio/Leica) connected to a web-based digital slide repository powered by eSlide Manager and housed in the Vanderbilt University Medical Center data center (*examples of islet images are shown in **Figures 1 and 2***).

Figure 1. Histological assessment of human islet composition. Human islet preparation was processed for cryosections. (A) Each specimen label is captured by slide scanner and contains information about UNOS ID, disease type, donor age, gender, and islet isolation center. (B - D). Islet composition was determined by immunofluorescence for islet markers (β cells – C-peptide, green; α cells – glucagon, red; δ cells – somatostatin; white, DAPI – blue) using a tissue classifier algorithm (HaloTM).

Figure 2. Histological assessment of human islet endocrine and acinar cell compartments. Human islet preparation was processed for cryosections. (A) Each specimen label is captured by slide scanner and contains information about UNOS ID, disease type, donor age, gender, and islet isolation center. (B - D) Sections were labeled for islet markers (C-peptide, glucagon, and somatostatin – green), acinar cell marker Hpx1 (red), counterstained with DAPI (blue) and then analyzed with algorithm (HaloTM) to measure endocrine cell compartment.

- 3.2** Using a tissue classifier algorithm (HaloTM, Indica Labs) analyze islet images (50 -100 islets/labeling experiment) to provide a quantitative assessment of the islet cell composition and endocrine/acinar cell compartments for a given human islet preparation (*examples of quantitative islet assessment are shown in the legend of **Figures 1 and 2***).

Data Storage and Reporting

4 Data Storage and Reporting

- 4.1** To facilitate data management and ensure data security, the Vanderbilt HIPP uses an institutional server-based platform for data storage and analysis.
- 4.2** Upon analysis completion (within 14 business days) annotated images containing metadata and image analysis outputs will be uploaded to the IIDP-HIPP database and immediately disseminated to IIDP-affiliated investigators and islet isolation centers.