DEC 18, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5bgg2jw

Protocol Citation: Integrated Islet Distribution Program 2019. Attachment 1: Preparation of Fluorescein Diacetate and Propidium Iodide Stock Solutions (FDA/PI). [protocols.io](https://dx.doi.org/10.17504/protocols.io.5bgg2jw) <https://dx.doi.org/10.17504/protocols.io.5bgg2jw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jul 09, 2019

Last Modified: Jun 11, 2020

PROTOCOL integer ID:
 25672

Attachment 1: Preparation of Fluorescein Diacetate and Propidium Iodide Stock Solutions (FDA/PI)

Integrated Islet Distribution Program¹

¹[Integrated Islet Distribution Program, City of Hope, Duarte, CA]

Integrated Islet Distribution Program

Tech. support email: iidp-email@coh.org

Integrated Islet Distribution Program

Integrated Islet Distribution Program, City of Hope

ABSTRACT

This Standard Operating Procedure is adapted from the work of the '*National Institutes of Health-Sponsored Clinical Islet Transplantation Consortium Phase 3 Trial: Manufacture of a Complex Cellular Product at Eight Processing Facilities*' following the SOP cited in the document '*Purified Human Pancreatic Islet - Viability Estimation of Islet Using Fluorescent Dyes (FDA/PI): Standard Operating Procedure of the NIH Clinical Islet Transplantation Consortium*'

This SOP defines the procedure for assessment of viability of human isolated islet preparations, which include endocrine and exocrine tissue, for use in the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) sponsored research in the Integrated Islet Distribution Program (IIDP). This protocol is written to assist the participating islet isolation centers and investigators who are part of this program.

Fluorescein Diacetate/ Propidium Iodide (FDA)/(PI) Viability Assay is a rapid fluorometric method to test the integrity of the plasma membrane simultaneously using inclusion and exclusion dyes; the assay differentiates between viable and nonviable cells and is, consequently, used for determination of viability of islet preparations.

Keywords: Inclusion and Exclusion Fluorescent Dyes, Fluorescein Diacetate, Propidium Iodide, Viability Assay

GUIDELINES

Fluorescein Diacetate/ Propidium Iodide (FDA)/(PI) Viability Assay is a rapid fluorometric method to test the integrity of the plasma membrane simultaneously using inclusion and exclusion dyes; the assay differentiates between viable and nonviable cells and is, consequently, used for determination of viability of islet preparations.

- The inclusion dye is Fluorescein Diacetate (FDA) and the exclusion dye is Propidium Iodide (PI). The final concentrations are as follows:
 - ◆ FDA: 0.46 μ M
 - ◆ PI: 14.34 μ M
- Fluorescein Diacetate is a nonpolar ester, which passes through plasma membranes and is hydrolyzed by intracellular esterases to produce free fluorescein. The polar fluorescein is confined within cells with an intact plasma membrane and can be observed under appropriate excitation conditions. FDA functions as an inclusion dye, i.e., viable cells will appear bright green fluorescent using FDA.
- Propidium iodide functions as an exclusion dye that cannot penetrate living cells but readily enters dead or dying cells. Once PI penetrates through the cell membrane, it binds to nucleic acids and causes them to fluoresce bright orange/red. PI absorbs in green light and fluoresces orange/red.
- Both of the fluorescent dyes used in this assay are light sensitive and must be kept in the dark, covered with aluminum foil.
- The fluorescent dyes are temperature sensitive and must be stored as follows:
 - ◆ FDA: $\leq -20^{\circ}\text{C}$
 - ◆ PI: $2-8^{\circ}\text{C}$

MATERIALS

MATERIALS

Fluorescein Diacetate Sigma Aldrich Catalog
#F7378

Propidium Iodide Sigma Aldrich Catalog
#P4170

Acetone Sigma Aldrich Catalog
#179124

Thermo Scientific™ Nunc™ Cell Culture/Petri Dishes Fisher Scientific Catalog
#174926

Equipment

OHAUS™ Explorer™ Semi-Micro Balance or equivalent

NAME

Semi-Micro Balance

TYPE

Ohaus

BRAND

01-919-370

SKU

<https://www.fishersci.com/shop/products/ohaus-explorer-semi-micro-balances/p-6541003#>

LINK

0.01 mg to 52 gm capacity

SPECIFICATIONS

Equipment

Snap Cap Microcentrifuge Tube or equivalent

NAME

Polypropylene Microcentrifuge Tube

TYPE

Corning Costar Snap Cap Microcentrifuge Tube

BRAND

07200210

SKU

<https://www.fishersci.com/shop/products/costar-microcentrifuge-tubes-6/07200210>

LINK

2 mL snap cap polypropylene micro tube

SPECIFICATIONS

Equipment	
5 mL Transport Tubes	NAME
5 mL Transport Tubes	TYPE
Globe Scientific Self-Standing Transport Tubes or	BRAND
22-010-1227	SKU
https://www.fishersci.com/shop/products/self-standing-transport-tubes-separate-screw-cap-7/p-7112488?keyword=5+ML+CENTRIFUGE+TUBE+WITH+CAP	LINK
5 mL Polypropylene tube with Polyethylene Screw cap	SPECIFICATIONS

Equipment	
Fisherbrand™ Pipette-Specific Tips	NAME
Pipette Tips, 100 to 1000uL or equivalent	TYPE
Pipet Tips	BRAND
02-681-182	SKU
https://www.fishersci.com/shop/products/fisherbrand-pipetter-specific-tips-natural-100-1000-l-101-6mm-long-bulk-pack/02681182#?keyword=true	LINK
100 to 1000uL	SPECIFICATIONS

Always wear gloves and observe standard chemical procedures:

Fluorescein Diacetate: [FDA MSDSAction.pdf](#)

- Protect from light. Avoid contact and inhalation. Nitrile gloves are recommended in the MSDS when handling FDA.

Propidium Iodide: [PI MSDSAction.pdf](#)

- Use personal protective equipment. Product may be toxic if inhaled, swallowed, or splashed on skin. Avoid dust formation. Avoid breathing vapors, mist, or gas. Ensure adequate ventilation. Wear gloves and observe Safety Data Sheet. Suspected of causing genetic defects.

Acetone: [AcetoneMSDSAction.pdf](#)

- Solvent/Flammable. Keep away from heat, spark, and open flame. Keep container tightly closed. Use with adequate ventilation. Avoid contact with eyes. Avoid prolonged or repeated breathing of vapor. Handle with gloves. Gloves must be inspected prior to use. Use proper glove removal technique (without touching glove's outer surface) to avoid skin contact with this product. Dispose of contaminated gloves after use in accordance with applicable laws and good laboratory practices. Wash and dry hands. Store aliquots in tightly sealed in glass or polypropylene tubes with polypropylene or polyethylene closure.

Preparation of Stock Fluorescein Diacetate (FDA) Solution

- 1 Prepare [\[M\]](#) 24 micromolar (μM) Fluorescein Diacetate (FDA) according to the formulation in sub-steps below.

Note

The expiration date, for both PI and FDA stains, is six months from the date of preparation.

Both of the fluorescent dyes used in this assay are light sensitive and must be kept in the dark, covered with aluminum foil.

1.1 Calculation:

FDA FW = 416.4

Stock Concentration = 24 μ M

Volume required = 200 mL

Formula: FW X Concentration X Volume = $(416.4) \times (24 \times 10^{-6}) \times (200 \times 10^{-3}) = 0.00199$ g FDA

Sigma Aldrich has a Mass Molarity Calculator that can be used to determine smaller quantities: <https://www.sigmaaldrich.com/chemistry/stockroom-reagents/learning-center/technical-library/mass-molarity-calculator.html>

1.2 Dissolve 0.00199 g of FDA in 200 mL of acetone in a glass bottle and cover with aluminum foil.

0.00199 g
FDA

- FDA Supplier _____
- FDA Lot # _____
- Expiration Date _____

200 mL
Acetone

- Acetone Supplier _____
- Acetone Lot # _____
- Expiration Date _____

1.3

- Store Stock FDA solution tightly sealed in 4 mL or smaller aliquots in polypropylene tubes or glass tubes at $\leq$ -20 °C , **protected from light**, for up to six months.

- Record the following on each aliquot tube:

FDA (24 μ M) Stock Solution Assigned Lot number: _____

Expiration Date: _____

Prepared By/Date: _____

Reviewed By/Date: _____

Preparation of Stock Propidium Iodide (PI)

- 2 Prepare 750 micromolar (μ M) Propidium Iodide (PI) according to the formulation in sub-steps below.

2.1 Calculation:

PI FW = 668.4

Stock Concentration = 750 μ M

Volume required = 25 mL

$$\text{FW} \times \text{Concentration} \times \text{Volume} = (668.4) \times (750 \times 10^{-6}) \times (25 \times 10^{-3}) = 0.0125 \text{ g PI}$$

2.2 Dissolve 0.0125 g of PI in 25 mL of DPBS and cover with aluminum foil.

0.0125 g Propidium
Iodide

- PI Supplier _____
- PI Lot # _____
- Expiration Date _____

25 mL
DPBS

- PBS Supplier _____
- PBS Lot # _____
- Expiration Date _____

2.3

- Store Stock PI solution tightly sealed in 0.5 mL or smaller aliquots in polypropylene Microcentrifuge Snap Cap tubes or glass tubes at 2 °C to 8 °C , **protected from light**, for up to six months.

- Record the following on each aliquot tube:

PI (750 μ M) Stock Solution Assigned Lot number: _____

Expiration Date: _____

Prepared By/Date: _____

Reviewed By/Date: _____