VERSION 3

JAN 14, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.8pihvke

Protocol Citation: Mariella Ferrante, Rossella Annunziata, Monia Teresa Russo, Francesco Manfellotto 2020. Axenic Diatoms cultures protocol. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.8pihvke>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 24, 2019**Last Modified:** Jan 14, 2020**PROTOCOL integer ID:**
29130

Axenic Diatoms cultures protocol V.3

Mariella Ferrante¹, Rossella Annunziata¹, Monia Teresa Russo¹,
Francesco Manfellotto¹

¹Stazione Zoologica Anton Dohrn

Francesco Manfellotto

Stazione Zoologica Anton Dohrn

ABSTRACT

Axenic cultures protocol

Axenicity of cultures is obtained by multi-antibiotic treatment: by adding antibiotics directly to the medium in which the strain will be inoculated. After a variable amount of days we refresh the coltures with new medium.

antibiotics concentration

- 1 The antibiotics used are streptomycin, penicillin and ampicillin;
We used the following final concentration:
 - Streptomycin 0,5 mg/ml
 - Penicillin 0,05 mg/ml
 - Ampicillin 0,03 mg/ml
- 2 Add antibiotics in F/2 medium.
Start from 20/50.000 cells in 25 ml of F/2 medium with antibiotics.
Refresh 3 times the cultures every 3 days with new medium with antibiotics.
- 3 Transfer the cultures into a variable higher volume flask containing F/2 medium and antibiotics.
Arrive progressively at the final volume.

contamination test

- 4 Assess axenicity of the strains by fluorescence microscopy using DAPI staining (4',6-diamidino-2-phenylindole-a DNA stain).

(DAPI stains DNA, if bacteria are present in the culture bacterial nucleoids can be visualised as fluorescent spots).

To fix cells use neutralized formaldehyde 1.6%
DAPI final concentration 1:1000

Pseudo-nitzschia multistriata fluorescence microscopy images using DAPI staining.

- A. Normal culture with bacteria
- B. Axenic culture without bacteria.