MAY 25, 2018

BACRB - Parallel Sequencing of Barcoded BAC Clones by random tagging PCR

Elena Hilario¹, Marcus Davy¹, Roy Storey¹, Richard Newcomb¹, Ross N Crowhurst¹, Roger P Hellens²

¹Plant and Food Research; ²Queensland University of Technology

Elena Hilario

Plant and Food Research

ABSTRACT

A genome is assembled by connecting large scaffolds via a collection of thousands of mate-paired ends of various sizes (3-20 kbp). Terminal ends of long DNA fragments can be obtained by sequencing large insert clones by Sanger sequencing, a labour-intensive and expensive approach. Here we describe a high-throughput protocol to tag 11,520 BAC plasmids randomly with barcoded oligonucleotides at the end of a random hexamer by PCR. This method assisted the kiwifruit (*Actinidia chinensis*) genome assembly in a similar way as mate paired end datasets. Although not fully sequenced, each randomly tagged BAC clone provides a collection of short sequence tags mapped to a genome across one or multiple scaffolds under three different scenarios: validate a single scaffold, join two scaffolds, or group several scaffolds by proximity only. Our method provides a simple way of cataloging BAC libraries to assist genome assembly projects.

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.p3adqie

External link:

<https://bmcbgenomics.biomedcentral.com/articles/10.1186/s12864-018-4656-3>

Protocol Citation:

Elena Hilario, Marcus Davy, Roy Storey, Richard Newcomb, Ross N Crowhurst, Roger P Hellens 2018. BACRB - Parallel Sequencing of Barcoded BAC Clones by random tagging PCR. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p3adqie>

MANUSCRIPT CITATION:

Pilkington SM, Crowhurst R, Hilario E, Nardoza S, Fraser L, Peng Y, et al. (2018) A manually annotated *Actinidia chinensis* var. *chinensis* (kiwifruit) genome highlights the challenges associated with draft genomes and gene prediction in plants. *BMC Genomics* 19: 257

License:

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status:

Working
We use this protocol and it's working

Created: May 11, 2018

Last Modified: May 25, 2018

PROTOCOL integer ID: 12098

Keywords: BAC sequencing, Illumina, random tagging PCR

GUIDELINES

A BAC library from kiwifruit (*Actinidia chinensis*) was constructed by Bio S&T, Québec, Canada. The nuclear genomic DNA was partially digested with *Hind*III, cloned in the plant transformable binary vector pCLD04541, maintained in *Escherichia coli* DHB10 and arrayed in 180 382-well plates. The BAC library represents ~ 6.7 times the haploid genome; the average insert size is 71.3 kbp (*n* = 309), and 0.6% of the library contain organellar DNA clones.

The process of oligonucleotide design will be reported in a separate protocol. But here is an overview:

Conditions for generating BAC Random Barcoded oligonucleotides (BACRB)

From all possible combinations ($4^{14} = 268,435,456$) select 384 unique barcodes with the following characteristics:

- GC = 0.5 ± 0.03
- No palindromes
- No repeats > 3 bases long
- Should not hybridize to plasmid vector or *Escherichia coli* genome
- 14 b long

An example...

Amplification step	Oligonucleotide set	Sequence
PEP-PCR	BACRB_Q1_A1	ACTTACACGTCGAGNNNNNN
TD-PCR	t_BACRB_Q1_A1	GGCGTA ACTTACACGTCGAG

SAFETY WARNINGS

Please read the safety warnings for the use of plate bench centrifuge and for bacterial and isopropanol waste disposal noted in the protocol.

BEFORE START INSTRUCTIONS

NOTES

1. When **pipetting dense solutions**, like the PCR master mix, prime the pipette tips by drawing and expelling the liquid slowly, one time. Multichannel pipettes (and robotic stations) allow you to select the speed of this operation. Also, if the required volume is 47.5 µL, set the pipette to 46.7 µL, and don't dip the tips too far into the solution. The amount of carry over solution outside the tip will compensate for the "missing" µL in the settings.
2. This method was developed in 2011/2012, and since then the library preparation and sequencing technologies have changed rapidly. You can select any approach as long as the amplicon ends are recovered during the library preparation. Do not use a tagmentation based protocol to prepare the libraries. And we recommend to obtain a pair end read data set to increase the read mapping rate.
3. We are aware that a small amount of *Escherichia coli* genomic DNA co-purifies with the BAC plasmid

and is tagged and amplified along with the target BAC DNA. The *E. coli* DNA can be digested with enzyme blends such as [PlasmidSafe](#) but given the number of BAC plasmid preps needed, it becomes too expensive. An option is to remove the contaminant by [genomic subtractive hybridization](#). A subtracter is prepared with purified *E. coli* DNA which is sheared, end repaired and biotinylated at the 3'-end with terminal deoxynucleotidyl transferase. The subtracter is hybridized with the plate pool amplicons and the double stranded *E. coli* molecules (subtracter+random tagged *E. coli* DNA) are removed with streptavidin magnetic beads. After 3 rounds of subtraction, the remaining amplicon sample should be fairly clean of host DNA and ready for library preparation.

Consumables

- ABgene 96 well polypropylene storage microplate, conical bottom (cat# AB1058)
- 2 mL deep well plate (NUNC cat# 278752)
- 96-well Filter plate (NUNC cat# 278011)
- Conical polypropylene sterile tubes, 15 mL
- Disposable sterile reservoirs
- Breathable tape (rayon acrylate NUNC cat# 241205)
- Silicone mat that fits the 2 mL deep well plate or a piece of silicone rubber sheet (1.5 mm thickness)
- Clear packaging tape (48 mm wide) for temporary plate sealing
- Cutter with retractable blade
- Microcentrifuges, 1.5 and 2 mL, sterile, and racks
- PCR plates (Thermo-Fast AB-0600)
- Aluminium tape for PCR reactions and long term storage (NUNC cat# 232698, or similar)
- Paper towels, low lint
- Rectangular plastic box
- Waste containers for biological material and isopropanol

Solutions

- [LB medium](#)
- 12.5 mg/mL tetracycline hydrochloride, dissolved in absolute ethanol
- RNase A 100 µg/µL (Qiagen cat# 19101)
- Resuspension buffer (25 mM Tris-HCl pH 8, 10 mM EDTA, sterile). Add RNase A to a final concentration of 50 µg/mL before resuspending the bacterial cells
- 0.2 N NaOH prepared in sterile deionized water and stored in a plastic bottle
- 10% SDS, sterile
- Lysis solution (Use sterile deionized water to prepare 0.2 M NaOH, 0.1% SDS. This solution is freshly made from stocks described above)
- Neutralizing solution (3 M potassium acetate, 2 M acetic acid, pH 5.6). To prepare this solution dissolve 295.5 g potassium acetate (MW 98.5 g/mol) in ~400 mL sterile deionized water. Adjust the volume to 600 mL and store in a sterile 1 L glass bottle. Add 115 mL glacial acetic acid and 285 mL sterile deionized water. Do not autoclave and do not adjust the pH, which should be around 5.6. Split the solutions in 2 or 3 sterile glass bottles and store at 4°C
- Isopropanol
- 70% ethanol, in sterile deionized water
- Plasmid resuspension buffer (28 mM Tris-HCl pH 8, 1 mM EDTA, 0.6 mM cresol red)
- Deionized water, sterile
- AccuPrime Taq DNA polymerase High Fidelity 5 U/µL and 10X AccuPrime High Fidelity Buffer I and II (Invitrogen 12346094)
- Mineral oil (Sigma-Aldrich cat # M5904-500ML)
- 2% xylene cyanole FF solution prepared in sterile deionized water
- Platinum *Pfx* DNA polymerase 2.5 U/µL (Invitrogen 11708039)
- 1.2 M Trehalose, dissolved in sterile deionized water and filtered through 0.2 or 0.4 µm

- 10X *Pfx* reaction buffer homemade: 200 mM Tris-HCl, 100 mM KCl, 100 mM (NH₄)₂SO₄, pH 8, sterile. **Note:** *I have not tried the 10X buffer and enhancer solutions supplied with the Platinum *Pfx* DNA polymerase (Invitrogen 11708039). It should work, but do a preliminary test*
- 50 mM MgSO₄ (use the stock provided with either DNA polymerase)
- 2 mM dNTP (dilution prepared in deionized sterile water from 100 mM stocks)
- 2% agarose gels in [1X TAE](#) buffer and 1X TAE running buffer
- 1 kb+ DNA ladder (Invitrogen) or similar
- SybrSafe (Invitrogen)
- 3 M sodium acetate pH 5.2. To prepare this solution dissolve 24.6 g sodium acetate anhydrous (MW 82.03 g/mol) in ~ 80 mL deionized water. Adjust the pH to 5.2 with glacial acetic acid. Adjust the volume to 100 mL, split into two glass bottles, and autoclave
- TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 7.5), sterile

Equipment and kits

- Cell culture incubator
- Bench top centrifuge with plate rotor (for example, Eppendorf SKU 5810000084 Rotor A-4-62, with 4 MTP buckets; Cat. # 5810711002)
- Bench top microcentrifuge for 1.5 mL tubes
- Vacuum manifold (for example Supleco PlatePrep 96-well Vacuum Manifold [SKU:575650-U](#))
- Vortex with plate adaptor
- Qubit 2.0 or above and High Sensitivity dsDNA kit
- Horizontal gel electrophoresis unit with power supply
- Image capture equipment (for example, GelDoc, LSA3000, etc)
- Multichannel pipettes (1-10 µL, 5-200 µL, 50-1000 µL) or liquid handling robotic station with appropriate multichannel tools and tips, for example Biomek series, Hamilton, Integra's Viaflo, etc.
- PCR machine
- Timer

BAC clones cultures

- 1 Transfer an aliquot of the BAC library glycerol stock stored in the 384-well library plate using the 96-pin HDRT of the robotic unit into the 96-well conical bottom working plate which contains 150 µL of LB medium, 12.5 µg/mL tetracycline per well.
Cover the plate with its corresponding lid and incubate at 37°C overnight.

Note

Thirty 384-well plates from the kiwifruit (*Actinidia chinensis*) BAC library were selected and represent 1X coverage of the genome (~750 Mbp). Each 384-well plate was divided in 4 quadrants to be transferred to 96-well working plates: A1 = quadrant 1; A2 = quadrant 2; B1 = quadrant 3; B2 = quadrant 4. A total of 120 quadrant plates were generated.
Here we describe how to process one 384-well library plate (4 quadrants) only. You can scale it up to the desired number of library plates according to the level of coverage required.

Note

Use a liquid handling robotic unit to manipulate the BAC library plates. The robot should include a 96-pin high density replicating tool (HDRT).

37 °C cell culture incubator

- 2 Transfer 20 µL of each well of the 96-well working plate to its corresponding address in a 2 mL 96-deep well plate filled with 1.2 mL LB medium and 12.5 µg/mL tetracycline per well. Seal the plate with breathable tape and incubate at 37°C in a cell culture shaker at 180 rpm for 16 h.

16:00:00

37 °C cell culture shaker incubator

Note

To increase the BAC plasmid yield, the plates can be stored in a cold room for ~ 6 h or overnight before harvesting the bacterial cells.

- 3 Collect the bacterial cells by centrifugation at 3000 rpm at room temperature for 25 min in a benchtop microtitre plate centrifuge. Remove the tape and pour out the medium by inverting the plate over a rectangular plastic box. Blot briefly over paper towels and seal with packaging tape.
Store the plates containing the bacterial cells at -20°C until all the plates have been processed.

Safety information

To dispose the used medium transfer it to a flask and autoclave at 121°C for 15 min. After sterilization the medium can be poured down the drain.

Safety information

Load the plates according to their matching weights. Use rectangular pieces of paper the same size as the plates to match the weight values and place it under the plates.

BAC plasmid preps

- 4 Remove the BAC cell culture plates from the freezer and let the pellets defrost at room temperature for 10-20 min.
- 5 Add 100 µL of resuspension buffer per well. Seal the plate with packaging tape and vortex for 10-20 seconds. Spin down briefly.

Note

The bacterial cell pellet should be completely dissolved.

- 6 Add 200 µL of lysis solution per well and gently swirl the plate to ensure complete lysis. Incubate without shaking for 5 min at room temperature.

⌚ 00:05:00

- 7 Neutralize the lysate by adding 150 μ L of ice cold potassium acetate solution per well. Seal the plate with packaging tape and gently mix by vortexing at the lowest speed, 3-4 pulses of 5 s each. Keep the plates on ice for 10 min.

⌚ 00:10:00

🧊 0 °C ice box

- 8 Centrifuge the plates for 15 min at 3000 rpm, room temperature.

⌚ 00:15:00 lysate centrifugation

Note

Most of the flocculent material is collected at the bottom of the well. Don't worry if some is transferred to the fritted plate in the next step.

⇒ go to step #3 undefined

- 9 Transfer the lysate supernatant to a NUNC fritted plate assembled over a vacuum manifold containing its corresponding 2 mL 96-deep well receiving plate, in the same orientation. Cover the fritted plate with a silicone mat and apply vacuum to -10 in Hg for 2 min or until all liquid has been filtrated. Discard the fritted plate.
- 10 Remove the receiving plate containing the BAC plasmid and add 0.7 volumes of isopropanol per well at room temperature (approximately 260 μ L) to each well. Seal the plate with packaging tape and vortex gently for 2-3 seconds, 3 times. Incubate at room temperature for 1 h.

⌚ 01:00:00

- 11 Collect the BAC plasmid DNA by centrifugation at 3000 rpm, 25 min at room temperature.

⌚ 00:25:00 BAC plasmid DNA

⇒ go to step #3 undefined

- 12 Pour out the supernatant into a rectangular waste tray. While the plate is still upside down, remove any traces of liquid by blotting it over paper towels for 2-3 seconds only.

Safety information

Dispose the isopropanol waste according to your local environmental regulations.

- 13 Add 100 or 200 μ L of 70% ethanol to each well. Seal the plate with packaging tape and swirl it gently. Let it stand at room temperature for 10-20 min. Collect the BAC DNA by centrifugation at 3000 rpm, 25 min at room temperature.

⇒ go to step #3 undefined

- 14 Pour out the ethanol supernatant over a rectangular tray and briefly blot the plate over paper towels. The ethanol waste can be poured down the drain and flushed with water for ~ 30 sec. Air dry the BAC pellet for ~ 10 min at room temperature. Add 150 μ L of plasmid resuspension buffer per well, seal with packaging tape, spin down briefly and store at 4°C.

Note

The purpose of adding cresol red to the BAC plasmid DNA solution is to ensure the pH of the DNA solution is >7.5, and to visually confirm that the template transfer was successful in the following PCR reaction. The BAC plasmid solution should be red/pink, not yellow, which means it still contains sodium acetate and the pH is below 7.5.

Note

Weight the plate at $t = 0$ and after 5 and 10 min or until you see no change in weight. But if you prefer not to over dry the BAC DNA prep, add the buffer, seal the plate with packaging tape, vortex at medium speed and spin down briefly. Remove the tape and let any remaining ethanol evaporate at room temperature for 10-20 min. Cover the plate with a tissue paper (lint free) to prevent foreign objects fall into the wells.

- 15 Quantify at least 12 BAC plasmid preps from each quadrant plate (48 BAC preps per library plate). Use 1 μL of BAC plasmid for the Qubit HS dsDNA kit. Take the average value of those 48 reactions and if needed adjust the concentration to $\sim 3 \text{ ng}/\mu\text{L}$ by adding more plasmid resuspension buffer.

Random tagging BAC clones - PEP-PCR

- 16 Make a master mix according to the following table, with at least 16 extra reactions:

PEP-PCR		
	One reaction, μL	400 reactions, μL
Deionised sterile water	38.8	15520
10X AccuPrime High Fidelity Buffer II	5	2000
BACRB oligonucleotide, 10 pmol/ μL	1	
AccuPrime Taq DNA polymerase High Fidelity 5 U/ μL	0.2	80
BAC plasmid prep ($\sim 3 \text{ ng}/\mu\text{L}$)	5	
Total volume	50	

Read note regarding pipetting dense solutions. Aliquot 44 μL of master mix per well into a PCR plate.

- 17 Add 1 μL of its corresponding BACRB oligonucleotide to each well. Add 5 μL of BAC plasmid solution to its corresponding well. Seal the plate with packaging tape and vortex vigorously. Spin down briefly. Add 20 μL of sterile mineral oil to each well with a multichannel pipette, seal the plate with aluminium tape.

Note

The addition of mineral oil should prevent evaporation but check all wells, especially the ones at the edge of the plate. If less than half of the total volume of the reaction has evaporated, add enough 1X TE pH 7.5 buffer to bring back the volume to 50 μL . Repeat the PEP-PCR reaction if more than half of the liquid has evaporated.

Avoid touching the rim of the well with mineral oil because it will prevent the aluminium tape stick properly.

- 18 Amplify as follows:

94°C, 2 min → (94°C, 40 sec → 30°C, 2 min → 48°C, 4 min (set ramp at +0.1°C/sec) → 68°C, 1 min) x 50 cycles → 68°C, 7 min → stop

and leave at room temperature (running time: 10 h)

Note

The annealing step is done at two temperatures: 30 and 48°C, with a very gentle ramp that will increase 0.1°C per second.

- 19** For short term storage, keep the plate at 4°C, otherwise store it at -20°C.

Note

The PEP-PCR plates are kept at 4°C until all the BAC plasmid preps have been processed. One microlitre of 2% xylene cyanole FF solution can be added to each PEP-PCR reaction before the next round of amplification to visually confirm the addition of template.

Enrich rt-BACs by touch down PCR

- 20** Make a master mix according to the following table, with at least 16 extra reactions:

TD-PCR		
	One reaction, μL	Final concentration
Sterile deionized water	0.7	280
1.2 M Trehalose	10	4000
10X <i>Pfx</i> Reaction Buffer (homemade)	4	1600
50 mM MgSO_4	0.2	80
2 mM dNTP	2	800
t-BACRB oligonucleotide, 4 pmol/ μL	1	4 pmol

Platinum <i>Pfx</i> , 2.5 U/μL	0 1	4 0	0. 2 5 u n i t s
PEP-PCR reaction, undiluted	2		
Total volume	2 0		

This is a dense solution. Aliquot 17 μL of master mix per well with a multichannel pipette into its corresponding PCR plate.

Note

No need to add BSA, it has no effect on the amplification and it only adds to the cost.
Platinum *Pfx* has been discontinued (April 2018) and replaced by another enzyme blend called Superfi (Invitrogen cat# 12351010). If you prefer to use another enzyme mix, just make sure is a proofreading type.

- 21 Add 1 μL of its corresponding t-BACRB oligonucleotide to each well. Add 2 μL of PEP-PCR solution to its corresponding well. Seal the plate with aluminium tape and vortex vigorously. Spin down briefly.
- 22 Amplify as follows:

94°C, 2 min → (94°C, 30 sec → Touchdown 68°C-58°C, 30 sec at -0.5°C/cycle → 68°C, 1 min) x 20 cycles → (94°C, 30 sec → 58°C, 30 sec → 68°C, 1 min) x 10 cycles → 68°C, 7 min → stop

and leave at room temperature.
- 23 Store the PCR plates at 4°C until all plates have been processed and you are ready to prepare the plate pools.

Prepare plate pools

- 24 Transfer the contents of each 96-well plate (one quadrant) into a 2 mL microcentrifuge tube (final volume ~ 1.8 mL) and analyze 20 μL by agarose gel electrophoresis (2% agarose in 1X TAE buffer, 1 kb+ ladder, 140 V, constant current, 30 min. Stain with SybrSafe 10 min (3 μL SybrSafe in 50 mL 1X TAE).

Expected result

- 25** Transfer 700 μ L of each quadrant pool into a 15 mL conical tube (final volume 2.8 mL). Add 280 μ L of 3 M sodium acetate pH 5.2 to each tube and vortex briefly. Add 1.96 mL isopropanol per tube at room temperature and vortex until completely mixed. Incubate at room temperature for 1 h.

🕒 01:00:00 isopropanol precipitation

⇒ go to step #12 undefined

- 26** Centrifuge the sample at 3000 rpm for 25 min. Discard the isopropanol supernatant and add 1 mL 70% ethanol per tube. Make sure the pellet is detached from the tube and keep at room temperature for ~ 10 min. Centrifuge again at the same speed and time.

⇒ go to step #3 undefined

- 27** Remove the ethanol supernatant and blot the tube on paper towels without losing the pellet. Air dry the pellet ~ 10 min and add 300 μ L TE buffer per tube.

⇒ go to step #14 undefined

- 28** Run 2 μ L of each plate pool in a 2% agarose gel in 1X TAE buffer.

⇒ go to step #24 undefined

Expected result

Illumina library prep and sequencing

- 29 The dataset reported in our paper was obtained by preparing TruSeq Illumina libraries. Each plate pool was fragmented, end repaired, dA tailed and barcoded with Illumina adaptors. Ten plate libraries were pooled per lane and sequenced in the HiSeq2000 platform, 100 cycles, single end mode. See 'Guidelines and Warnings' section for comments on library and sequencing platform improvements developed since this method was created.