

Oct 22, 2019

Bacterial Transformation

DOI

dx.doi.org/10.17504/protocols.io.8irhud6

Dana Mozaffari¹, Laura Kvedarauskaite¹

¹EPFL - EPF Lausanne

iGEM EPFL



Dana Mozaffari

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.8irhud6

External link: <https://international.neb.com/protocols/0001/01/01/high-efficiency-transformation-protocol-c3019>

Protocol Citation: Dana Mozaffari, Laura Kvedarauskaite 2019. Bacterial Transformation. protocols.io

<https://dx.doi.org/10.17504/protocols.io.8irhud6>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 21, 2019

Last Modified: October 22, 2019

Protocol Integer ID: 28977

Abstract

This protocol explains how to transform a plasmid DNA template into competent bacterial cells which are then cultivated. The DNA is then purified from the cells using a miniprep kit.

The competent cells used in this protocol come from the NEB[®]10-beta Competent *E. coli* kit. This protocol was adapted from the one provided with the cells.

Materials

Materials required

- a tube of NEB 10-beta Competent *E. coli* cells
- 1-5 μ l containing 1 pg-100 ng plasmid DNA template
- selection plates with appropriate antibiotic
- DNA miniprep kit



DNA Transformation

- 1 Thaw a tube of NEB 10-beta Competent *E. coli* cells on ice for 10 minutes
- 2 Add 1-5 μ l containing 1 pg-100 ng of plasmid DNA to the cell mixture. Flick the tube 4-5 times to mix cells and DNA but do not vortex.
- 3 Place the mixture on ice for 30 minutes. Do not mix.
- 4 Heat shock at exactly 42°C for exactly 30 seconds. Do not mix.
- 5 Place on ice for 5 minutes. Do not mix.
- 6 Pipette 950 μ l of room temperature NEB 10-beta/Stable Outgrowth Medium into the mixture.
- 7 Place at 37°C for 60 minutes. Shake vigorously (250 rpm) or rotate.
- 8 Warm selection plates to 37°C.
- 9 Spread 50-100 μ l of each dilution onto a selection plate and incubate overnight at 37°C. Alternatively, incubate at 30°C for 24-36 hours or 25°C for 48 hours.

DNA Miniprep

- 10 Purify the DNA from selected cultures using a DNA miniprep kit