

Jun 21, 2019

Barcode plasmid library cloning

DOI

dx.doi.org/10.17504/protocols.io.4hgg3w

Benjamin Emert¹

¹University of Pennsylvania

RajLab



Benjamin Emert

University of Pennsylvania

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

[Create free account](#)

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.4hgg3w>

Protocol Citation: Benjamin Emert 2019. Barcode plasmid library cloning. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4hgg3w>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development

We are still developing and optimizing this protocol

Created: June 21, 2019

Last Modified: April 05, 2024

Protocol Integer ID: 24840

Keywords: cloning time machine barcode plasmid library, time machine barcode plasmid library, barcode plasmid library, cloning protocol

Abstract

Protocol for cloning Time Machine barcode plasmid library.

Materials

MATERIALS

-  Gibson Assembly Master Mix - 10 rxns **New England Biolabs Catalog #E2611S**
-  Electroporation Cuvette 1mm **Bio-Rad Laboratories Catalog #1652089**
-  Monarch® PCR & DNA Cleanup Kit (5 µg) **New England Biolabs Catalog #T1030**
-  LB plates with 100 µg/ml ampicillin
-  LB Broth with 100ug/mL ampicillin **Gibco - Thermo Fisher Scientific Catalog #244620**
-  Lucigen Endura electrocompetent cells **Lucigen Catalog #60242**
-  ssDNA barcode template **IDT**
-  Linearized vector backbone
-  EndoFree Plasmid maxiprep kit **Qiagen Catalog #12362**

Troubleshooting

Gibson assembly

- 1 Perform Gibson assembly using 100ng linearized vector and 5:1 molar excess of ssDNA template per 20 microliter reaction. 
- 1.1 Pre-heat thermal cycler to 50°C with lid set to 65°C.
 -  50 °C Pre-heat thermal cycler
- 1.2 Combine 100ng linearized vector with ~21.7 fmol ssDNA template (1.08 microliters of 100nM oligo) to a final volume of 10 microliters in nuclease free water. In parallel, perform a negative control reaction with plasmid and nuclease free water without the ssDNA insert. Keep on ice.
 -  100 ng Linearized vector
 -  1.08 µL ssDNA oligo (160nt)
 -  10 µL Final volume in NF water
- 1.3 Thaw Gibson assembly master mix on ice then add 10 microliters to plasmid plus insert. Pipette mix then spin down if needed.
 -  10 µL Gibson Assembly master mix
- 1.4 Incubate on thermal cycler at 50°C for 1 hour.
 -  50 °C
 -  01:00:00 incubation
- 2 Purify Gibson reaction using PCR clean-up column according to manufacturer's protocol eluting in 20 microliters of
 - 2.1 Add 40 microliters of DNA binding buffer to 20 microliters of Gibson assembly.
 -  40 µL DNA binding buffer
 -  20 µL Gibson assembly
 - 2.2 Re-load eluant and re-spin.

2.3 Add 200 microliters DNA wash buffer. Make sure ethanol has been added to wash buffer before use.

 200 μ L Wash buffer

2.4 Centrifuge at 12,000 x g for 1 minute.

 12000 x g

 00:01:00

2.5 Repeat wash and centrifugation (steps 2.4 and 2.5)

2.6 Transfer column to a new eppendorf or collection tube then centrifuge at 12,000 x g for 2 minutes to dry column

 12000 x g

 00:02:00

Note

Sometime residual liquid can collect on a lip inside the column. It may be helpful to rotate the column 180° after a 1 minute spin, then repeat another 1 minute spin to get rid of this liquid.

2.7 Add 10 microliter of nuclease free water pre-heated to 50°C to column and incubate for \geq 2 minutes

 10 μ L nuclease free water

 50 °C Pre-heat water

 00:02:00 Incubate

2.8 Centrifuge at 12,000 x g for 1 minute to elute

 12000 x g

 00:01:00

2.9 Optionally nanodrop the recovered DNA. Expected concentration is 10-15ng/uL.

Electroporation

- 3 Electroporation of competent cells
 - 3.1 Pre-warm Lucigen recovery media to 37°C

 37 °C Pre-warm recovery media
 - 3.2 If using Biorad Gene Pulser, pre-chill cuvettes to 4°C.

 4 °C Pre-chill electroporation cuvettes
 - 3.3 Thaw Lucigen Endura electrocompetent cells on ice. Each DUO vials contains 50uL of cells, enough for 2 electroporations. Use one reaction's worth of cells to electroporate the negative control Gibson assembly.
 - 3.4 Pipet vial of electrocompetent cells into 2 25 microliter aliquots.
 - 3.5 Add 2 microliters of cleaned Gibson assembly to 25 microliters of electrocompetent cells.

 2 µL Gibson assembly per  25 µL electrocompetent cells
 - 3.6 Transfer electrocompetent cells to pre-chilled cuvette. Keep on ice.
 - 3.7 Using the GenePulserXCell, electroporate with the following manual pulse setting: PL - 25msec, C - 10uF, PC- 600Ohms, V-1800

Note

Pulse length - 25msec, Capacitance - 10 microFaraday, Resistance - 600 ohms, Voltage - 1800 Volts.

Note

Expected time constant around 5-6 msec.

3.8 Immediately add 975 microliters of pre-warmed recovery media to cuvette then transfer to clean eppendorf or vial.

3.9 Incubate on shaker at 250 rpm at 37°C for 1 hour.

 37 °C  01:00:00 On shaker at 250rpm

3.10 Plate dilutions of electroporated cells for plating and calculating electroporation efficiency.

Prepare first dilution by adding  10 µL electroporated cells to  990 µL LB Broth

Prepare second dilution by adding  100 µL First dilution to  900 µL LB broth

Plate  100 µL first dilution on pre-warmed LB/ampicillin agar plate for a 1:1000 dilution plate.

Plate  100 µL second dilution on pre-warmed LB/Ampicillin agar plate for a 1:10,000 dilution plate.

Expected result

After incubating the plates overnight, expect to see >1,000 colonies on the 1:1000 dilution plate and >200 colonies on the 1:10,000 plate.

3.11 Add the remainder of the electroporated cells to 150mL LB broth containing 100ug/mL ampicillin. Incubate 12-14 hours on shaker at 32°C.

 150 mL LB broth

 32 °C

 12:00:00 -  14:00:00 shaker

3.12 Pellet the overnight culture by centrifugation in 50mL falcon tubes then store pellets at -20°C or proceed immediately with maxiprep.

 6000 x g  00:15:00  4 °C Pellet e. coli cultures.

 -20 °C Store e. coli pellets

Note

If centrifuge capable of 6,000 x g is unavailable, can pellet at $\geq 3,000$ x g.

4 PCR screen colonies to confirm barcode insertion.

5 Maxiprep barcode plasmid library.