VERSION 1
OCT 14, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bmatk2en

Protocol Citation: Matthew Haines 2020. BHI + v2 salts media. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bmatk2en>

MANUSCRIPT CITATION:
Weinstock, M. T., Hessek, E. D., Wilson, C. M., & Gibson, D. G. (2016). *Vibrio natriegens* as a fast-growing host for molecular biology. *Nature Methods*, 13(10), 849–851.
<https://doi.org/10.1038/nmeth.3970>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Sep 11, 2020

BHI + v2 salts media V.1

Matthew Haines¹

¹Imperial College London

Matthew Haines
Imperial College London

ABSTRACT

Vibrio natriegens grows exceptionally well in BHI + v2 salts media ([link](#)). Furthermore, this media has been used multiple times in the literature for culturing this organism. However, care must be taken when preparing this media not to autoclave v2 salts and BHI together. This protocol utilises a 10x v2 salts buffer to achieve this.

MATERIALS

MATERIALS

MgCl₂ **Contributed by users**

Sodium chloride **Contributed by users**

Potassium Chloride **Contributed by users**

Brain Heart Infusion Broth Dry Medium **Teknova Catalog #B9500**

Prepare 10x v2 salts buffer

- 1 Dissolve the following salts in 800 mL of ddH₂O:

Component	MW	Target concentration (mM)	Component (g/900 mL)	Final concentration (mM)
NaCl	58.44	2040.0	107.30	2040.08
MgCl ₂	95.21	231.4	19.83	231.42
KCl	74.55	42.0	2.82	42.03

- 2 Adjust the volume to 900 mL using ddH₂O.

Prepare BHI media

- 3 Dissolve 37 g of BHI dry medium in 900 mL ddH₂O.

Sterilise and combine

- 4 Sterilise both BHI media and 10x v2 salts buffer by autoclaving.
- 5 Add 100 mL 10x v2 salts buffer to the sterilised 900 mL BHI media under sterile conditions.

Note

The remaining 10x v2 salts buffer can be used for making further BHI + v2 salts media.