



Aug 15, 2020

Binding properties of Human IgG to immunoglobulin-binding proteins tested by double immunodiffusion (Ouchterlony) technique.

DOI

dx.doi.org/10.17504/protocols.io.bjskncw

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West In...

angel.vaillant@sta.uwi.e...



Angel A Justiz-Vaillant

University of the West Indies St. Augustine

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.bjskncw>

Protocol Citation: Angel A Justiz-Vaillant 2020. Binding properties of Human IgG to immunoglobulin-binding proteins tested by double immunodiffusion (Ouchterlony) technique.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjskncw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 15, 2020

Last Modified: August 15, 2020

Protocol Integer ID: 40492

Keywords: montage antibody purification kit with prosep, properties of human igg, antibody purification kit, montage antibody purification kit, immunoglobulin, human sera by affinity chromatography, human igg, affinity chromatography, binding protein, double immunodiffusion, purification, prepared in the laboratory, laboratory by the periodate method

Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK




The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://www.protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://www.protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

1. IBP-HRP conjugates can be prepared in the laboratory by the periodate method.
2. The Human IgG used can be commercially available or prepared in the laboratory from human sera by affinity chromatography (using an SpG-agarose column).
3. SpG-agarose column are widely commercially prepared.
4. PURE-1A antibody purification kit will be discontinued. However there are others product for this purpose as Montage Antibody Purification Kit with PROSEP-G media.

Materials

MATERIALS

-  Affinity Purification Using HiTrap™ Columns
-  Montage Antibody Purification Kit with PROSEP-G media
-  PURE-1A (antibody purification kit)-- Sigma Aldrich

Troubleshooting

- 1 The binding of Human IgG to peroxidase-conjugated Immunoglobulin-binding protein (IBP-HRP) including SpA-HRP, SpG-HRP, SpL-HRP, SpLA-HRP, SpAG-HRP, SpLG-HRP, SpLAG-HRP and turtle serum (Sigma-Aldrich) as a negative control are investigated by double immunodiffusion.
- 2 Briefly, 1% agarose gels are prepared and wells cut into the gel using a template.
- 3 Initially, aliquots of 25 μ l of purified human IgG (Sigma-Aldrich) at a concentration of one (1) μ g/ μ l are applied to the centre well. Human IgG could be prepared homemade by purifying human serum by SpA-affinity chromatography (PURE-1A, Sigma-Aldrich).
- 4 The peripheral wells are filled with 25 μ l each of bacterial protein conjugates including SpA-HRP, SpG-HRP, SpL-HRP, SpLA-HRP, SpAG-HRP, SpLG-HRP, SpLAG-HRP diluted 1:5.
- 5 The gels are incubated at RT for 48–72 hours incubated in a humidity chamber.
- 6 After that the gels are examined for precipitin lines.
- 7 Turtle serum is included as negative controls and all IBP-HRP conjugates suppose to react with human IgG.
- 8 The positive results are taken as the presence of precipitin line/s and negative results, the absence of precipitin lines.