AUG 15, 2020

🌐 Binding properties of Staphylococcal protein A (SpA) to purified immunoglobulins tested by double immunodiffusion (Ouchterlony) technique.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjsnknde

Protocol Citation: Angel A Justiz-Vaillant 2020. Binding properties of Staphylococcal protein A (SpA) to purified immunoglobulins tested by double immunodiffusion (Ouchterlony) technique..

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bjsnknde>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 15, 2020

Last Modified: Aug 15, 2020

PROTOCOL integer ID:
40494

1 The binding of SpA to purified IgG are investigated by double immunodiffusion.

- 2 Briefly, 1% agarose gels are prepared and wells cut into the gel using a template.
- 3 Initially, aliquots of 25 µl each of SpA diluted 1:5 are applied to the centre well.
- 4 The peripheral wells are filled with 25 µl each of purified IgG (1 µg/µl) in PBS pH 7.4.
- 5 The gels are incubated at RT for 48–72 hours.
- 6 After that the gels are examined for precipitin lines.
- 7 Human IgG are included as positive control and turtle serum as a negative control.
- 8 The positive results are taken as the presence of precipitin line/s and negative results, the absence of precipitin lines.