AUG 14, 2019

Biochemical Measures of Neuropathy - DHE

Eva Feldman¹

¹University of Michigan - Ann Arbor

Diabetic Complications Consortium

Tech. support email: rmcindoe@augusta.edu

Lili Liang

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.3n8gmhw

External link:

<https://www.diacomp.org/shared/document.aspx?id=54&docType=Protocol>

Protocol Citation:

Eva Feldman 2019. Biochemical Measures of Neuropathy - DHE. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3n8gmhw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status:

Working
We use this protocol and it's working

Created: Jun 03, 2019

Last Modified: Aug 14, 2019

PROTOCOL integer ID:

24000

ABSTRACT

Summary:

Oxidative stress is highly correlated with the metabolic changes caused by hyperglycemia. Increased levels of glucose overload mitochondria and result in the production of reactive oxygen species (ROS). In addition, the flow of excess glucose through cellular pathways decreases the cell's normal ability to detoxify ROS. As a result, the neurons and axons of the peripheral nervous system contain increased levels of ROS and decreased antioxidant capacity. The following assays are used to measure these changes in rodent models of diabetic neuropathy.

Diabetic Complication:

Neuropathy

Keywords: Biochemical Measures of Neuropathy, diabetic neuropathy, DHE

MATERIALS

MATERIALS

⊗ 10 mM HEPES **Molecular Probes**

⊗ 150 mM NaCl **Molecular Probes**

⊗ 5 mM KCl **Molecular Probes**

⊗ 1 mM MgCl₂ **Molecular Probes**

⊗ 1.8 mM CaCl₂ **Molecular Probes**

⊗ DHE **Molecular Probes**

⊗ DMSO (10 mg/mL) **Molecular Probes**

Reagent Preparation:

HBSS (pH 7.4): Combine HEPES, NaCl, KCl, MgCl₂, and CaCl₂. pH to 7.4 then filter to sterilize. (may be stored at 22°C for several months)

DHE: Dissolve 1 vial in 100 µL DMSO (10 mg/mL). Remove 10 µL, aliquot and freeze remainder. Protect from light and discard after thawing 3 times. Add the 10 µL to 1 mL HBSS (0.1 mg/mL), then for the working dilution put 100 µL of diluted solution into 10 mL HBSS. (1 µg/mL=3µM)

Note:

Molecular Probes ([RRID:SCR_013318](#))

1 Open Fluoroskan and choose open under the file menu. Scroll down and select DHE.sed. Set up your plate layout. Save your layout as DHxxxxxx.sed with xxxxxx being yy/mm/dd.

2 *Treat cells per experimental paradigm.

Note: *As an alternative, cells can be pre-loaded with DHE- do steps 3, 4, 5, 2, then 5.

3 15 minutes prior to reading, **gently** rinse cells once with HBSS.

4 Apply 3 µMDHE in HBSS and leave on for 15 minutes.

- 5 Rinse cells and add HBSS.
- 6 Place plate into Fluroskan holder and click **START**.
- 7 Take readings using 485 nm ex, 612 nm em filter pairs for ethidium and 355 nm ex, 430 nm em for DHE.
- 8 Save **both** sheets with .xls extension into the **DHE** data folder or your own folder. Use the naming convention **DHXXXX.xls**, where XXXX is the date in mmdd format and add a **r** for the red reading and **b** for the blue reading.