APR 26, 2019

Biolistic transformation of polar diatom *Fragilariopsis cylindrus*

Amanda Hopes¹, Thomas Mock¹¹University of East Anglia

Amanda Hopes

ABSTRACT

Transformation system for the polar diatom *Fragilariopsis cylindrus* using microparticle bombardment.

ATTACHMENTS

[FC transformation of IO protocols.docx](#)

GUIDELINES

Keep cells and plates at 4°C where possible throughout the protocol. Where brief room temperature steps are required, such as shooting cells, keep plates on ice before and transfer back to ice/4°C as quickly as possible.

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.z39f8r6

External link:

<https://ueaeprints.uea.ac.uk/66542/>

Protocol Citation: Amanda Hopes, Thomas Mock 2019. Biolistic transformation of polar diatom *Fragilariopsis cylindrus*. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.z39f8r6>

MANUSCRIPT CITATION:

Hopes A (2017). Expanding the molecular toolbox in diatoms: developing a transformation system, CRISPR-Cas and Inverse Yeast-1-hybrid. PhD Thesis, University of East Anglia.

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Apr 16, 2019

Last Modified: Apr 26, 2019

PROTOCOL integer ID:
22369

Keywords: Transformation,
Biolistic, Microparticle
bombardment, Diatom,
Fragilariopsis cylindrus, polar,
psychrophilic, psychrophile,
Antarctic

- 1 Culturing *Fragilariopsis cylindrus*. Grow *Fragilariopsis cylindrus* in Aquil media + Si (Price, 1989; <https://ncma.bigelow.org/algal-recipes>) at 4°C in 24 hour light. When culturing pre-chill the media and keep *F. cylindrus* on ice. Inoculate cells at a starting concentration of 50,000 cells/ml.
- 2 Making plates for shooting and selection. Make 1.5% Aquil agar plates for shooting and 0.8% Aquil agar plates supplemented with 100µg/ml zeocin (final concentration) for selection. To make plates make 2x concentrated SOW and 2x concentrated agar in separate bottles. Autoclave, cool to 50°C and mix. Add in Aquil nutrients once the SOW and agar have been mixed. For selective plates add in zeocin (stock concentration 100mg/ml). Pour plates and store at 4°C once set. Make sure plates are chilled to 4°C before using.
- 3 Plasmid. A plasmid containing a FCP:shble cassette and a FCP:egfp cassette have been used to transform *F. cylindrus*. Shble confers resistance to the antibiotic zeocin and the egfp has a human codon bias. An endogenous FCP promoter and terminator have been used. The plasmid can be found at Addgene: <https://www.addgene.org/85987/>
- 4 Preparation for shooting. Preparation of particles for shooting is based on the method by Kroth (2007). Weigh 60mg of M10 (0.7µm diameter) tungsten particles (BioRad, Hercules, CA) into a sterile 1.5 mL reaction tube. Add 1ml 100% ethanol and vortex for 3-5 minutes. Centrifuge at 12000 rpm for 1 minute and discard supernatant. Add 1ml of sterile water, resuspend, centrifuge for 1 minute at 12000 rpm and discard supernatant. Repeat two more times. Resuspend in 1ml of water and pipette 50ul aliquots into fresh Eppendorf tubes. To avoid sedimentation of particles during this step vortex frequently. Store aliquots at -20°C for up to two weeks. Make a 2.5M CaCl₂ solution (store at -20°C) and a 0.1M Spermidine solution (store at -20°C and use within one month).
- 5 Harvest cells. Harvest cells during exponential growth (5x10⁵ – 1x10⁶ cells/ml). Use 5 x 10⁷ cells for each shot. Filter cells at 4°C onto a 47 mm diameter 1.2 µm isopore filter (Merck, catalog number: RTTP04700) using vacuum filtration. Place the filter into the centre of a pre-chilled 1.5% agar Aquil plate. Keep plates at 4°C until needed for shooting, at which point transfer to ice.
- 6 Coat particles for shooting with the plasmid. Particles are coated according to Kroth (2007). Thaw

aliquots of CaCl₂, spermidine and tungsten particles. Resuspend tungsten particles by briefly vortexing. For 5 shots, pipette 50µl of tungsten particles, 5µg of plasmid (in 5-10µl of water), 50µl of 2.5M CaCl₂, and 20µl of 0.1M spermidine into a 1.5ml Eppendorf tube. Vortex for 1 minute. Centrifuge briefly and remove supernatant. Add 250µl of 100% ethanol and vortex until homogenous. Centrifuge briefly and remove supernatant. Add 50µl of 100% ethanol and vortex. Store on ice and use particles within 1 hour. Sterilise macrocarriers in 70% ethanol and leave to dry. Place into a sterilised macrocarrier holder (sterilised by 70% ethanol or autoclave), vortex prepared particles to resuspend and pipette 10µl into the centre of the macrocarrier. Leave to dry (make sure the ventilation hood is switched off to prevent particle migration) – this only takes a few minutes and proceed to shooting.

- 7 Microparticle bombardment. A PDS-1000/He biolistic microparticle delivery system (Bio-Rad Laboratories, catalog number: 1652257) was used to transform *F. cylindrus*. Prior to shooting autoclave the rupture disc retaining cap and microcarrier launch assembly. Wipe the particle gun inside and out with 70% ethanol. Set-up the device according to manufacturer's instructions. Dip a rupture disc (1550 psi) into 100% isopropanol, leave to dry and assemble into the rupture disc retaining cap. Clean helium lines by firing a shot without cells and particles. Perform particle bombardment as follows: Load a rupture disc into the retaining cap and install into the assembly. Load a stopping screen and the prepared macrocarrier holder into the microcarrier launch assembly. Place the plate with the chilled cells into the chamber (at room temperature) at a flight distance of 6cm. Create a vacuum of 25 Hg in the biolistic chamber, shoot and immediately return the cells to ice. Turn the filter paper upside down to bring the cells into contact with the Aquil plate surface and incubate at 4°C for 24 hours under standard conditions to allow cells to recover before selection. Shots are carried out in triplicate.
- 8 Selection. Keeping the plate on ice, rinse cells from the plate/filter with 500µl – 1ml of chilled Aquil media. Divide the cells between 5 selective plates on ice and spread with a sterilised drigalski spatula. Incubate cells at 4°C under standard conditions. Colonies typically take 3-7 weeks to appear at which point they can be screened by colony PCR or transferred to liquid media for further growth.
- 9 References. Price, N.M., Harrison, G.I., Hering, J.G., Hudson, R.J., Nirel, P.M., Palenik, B. and Morel, F.M., 1989. Preparation and chemistry of the artificial algal culture medium Aquil. *Biological oceanography*, 6(5-6), pp.443-461. Kroth, P.G., 2007. Genetic transformation; a tool to study protein targeting in diatoms, chap. 17. *Methods in molecular biology*, 2nd edn., Totowa, NJ, USA: Humana Press.