NOV 06, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.xiufkew

Protocol Citation: Dennis DD Dienst, Oliver Mantovani, Pia Lindberg 2019. (-)-α-Bisabolol GC sample preparation.

protocols.io
<https://dx.doi.org/10.17504/protocols.io.xiufkew>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jan 29, 2019

Last Modified: Nov 06, 2019

PROTOCOL integer ID:
 19764

Keywords: GC, bisabolol, dodecane, gas chromatography, terpenoids

(-)-α-Bisabolol GC sample preparation

Forked from [\(E\)-α-bisabolene GC sample preparation](#)

Dennis DD Dienst¹, Oliver Mantovani², Pia Lindberg¹

¹Department of Chemistry - Microbial Chemistry, Ångström Laboratory, Uppsala, Sweden;

²Universität Rostock, Germany

CyanoWorld

Metabolomics Protocols & Workflows

[1 more workspace](#) ↓

Dennis DD Dienst

Department of Chemistry - Ångström, Uppsala

ABSTRACT

This is a quick guide for the preparation of (-)-α-bisabolol samples and external standards in dodecane for GC analysis. This protocol has been established in the Lindberg lab at Ångström laboratory (Uppsala University) for direct analysis of dodecane-based *ex-situ* extracts from cyanobacterial strains producing (-)-α-bisabolol.

The protocol includes the preparation of stock solutions for the internal standard β-caryophyllene (BCP) and the external standard. The preparation of a fresh external standard series is suggested for each sample analysis. The calibration curve is required for quantification of (-)-α-bisabolol in the samples of interest.

This protocol is used in combination with the following GC protocol:

dx.doi.org/10.17504/protocols.io.kj2cuqe

MATERIALS

MATERIALS

β-Caryophyllene ≥80%, FCC, FG **Sigma Aldrich Catalog #W225207**

HPLC/GC Vials 1.5 mL clear glass **Vwr Catalog #548-1488**

Dodecane Reagent Grade ≥99% **Contributed by users Catalog #D221104**

Patchouli alcohol primary reference standard **Sigma Aldrich Catalog #5986-55-0**

Preparation of BCP (β -caryophyllene) internal standard (IS)..

1

		BCP Standard		
	Stock A	1: 10 Dilution from Original BCP Stock ($\Rightarrow 89 \text{ mg} \cdot \text{mL}^{-1}$)		
		↓		
	Stock B	281 μL Stock A + 719 μL dodecane ($\Rightarrow 25 \text{ mg} \cdot \text{mL}^{-1}$)	$\Rightarrow$	1: 100 to samples (2 μL + 198 μL sample)
		↓		
	Stock C	200 μL Stock B + 19,8 mL Dodecane ($\Rightarrow 250 \text{ } \mu\text{g} \cdot \text{mL}^{-1}$)	$\Rightarrow$	use for preparation of α -bisabolene external standard (ES) series

Store Stocks in the fridge at $\sim 4^\circ\text{C}$

Safety information

Dodecane is toxic! Wear protective gloves and goggles. Work under the fume hood, or use a respirator!

<https://pubchem.ncbi.nlm.nih.gov/compound/dodecane#section=Handling-and-Storage>

Preparation of (-)- α -Bisabolol standard stocks

2

	(-)- α - bisabolol standard
Stock L-I 200 μ L	1:100 dilution from original Stock ($\Rightarrow$ 9.2 μ g/ μ L in Stock C)
	$\Downarrow$
Stock L-II 2 mL	174 μ L Stock L-I + 1826 μ L Stock C ($\Rightarrow$ 800 μ g * mL ⁻¹)

Safety information

Dodecane is toxic! Wear protective gloves and goggles. Work under the fume hood, or use a respirator!

<https://pubchem.ncbi.nlm.nih.gov/compound/dodecane#section=Handling-and-Storage>

Preparation of (-)- α -Bisabolol calibration curve

3

Example: (-)- α -bisabolol external standard (ES) dilution series:

0; 12.5; 25; 50; 100; 200; 400; 800 μ g * μ L⁻¹

High Range Dilution Series:			

	Vol. S tock L-II	Vol. Stock C	Trans fer to vial (3 x)
800 $\mu\text{g} \cdot \text{mL}^{-1}$	600	0	200
400 $\mu\text{g} \cdot \text{mL}^{-1}$	325	325	200
200 $\mu\text{g} \cdot \text{mL}^{-1}$	162,5	487,5	200
100 $\mu\text{g} \cdot \text{mL}^{-1}$	81,25	568,75	200
50 $\mu\text{g} \cdot \text{mL}^{-1}$	40,625	600	200
25 $\mu\text{g} \cdot \text{mL}^{-1}$	20,3125	629,6875	200
12.5 $\mu\text{g} \cdot \text{mL}^{-1}$	10,15625	639,84375	200
0 $\mu\text{g} \cdot \text{mL}^{-1}$	0,0	600	200
Sum	1229,6875	2610,9375	

Safety information

Dodecane is toxic! Wear protective gloves and goggles. Work under the fume hood, or use a respirator!

<https://pubchem.ncbi.nlm.nih.gov/compound/dodecane#section=Handling-and-Storage>

Preparation of biological (-)- α -Bisabolol samples (in dodeca..

- 4
  - pipette 198 μL sample to GC vial
  - add each 2 μL **Stock B**

Safety information

Dodecane is toxic! Wear protective gloves and goggles. Work under the fume hood, or use a respirator!

<https://pubchem.ncbi.nlm.nih.gov/compound/dodecane#section=Handling-and-Storage>