AUG 30, 2018

Bodo saltans SSU-GFP-Neo cassette

Virginia

Fatma Gomaa¹, Edgcomb²,

Roberto

Docampo³

Peter Girguis¹, Zhuhong li³,

¹Harvard University; ²Woods Hole Oceanographic Institution;

³University of Georgia

Protist Research to Optimize Tools in Genetics (PROT-G)

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.s5meg46

Document Citation: Fatma Gomaa, Virginia Edgcomb, Peter Girguis, Zhuhong li, Roberto Docampo 2018. Bodo saltans SSU-GFP-Neo cassette. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.s5meg46>

License: This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: Aug 30, 2018

Last Modified: Aug 30, 2018

DOCUMENT integer ID:
15245

Fatma Gomaa

Harvard University

ABSTRACT

This plasmid is designed to integrate within the small subunit ribosomal RNA gene region of *B. saltans*. It carries 1 kb of the *B. saltans* small subunit ribosomal RNA gene. To construct this cassette, we first sequenced the complete ribosomal operon of *B. saltans*, which has a total length of ~9 kb. The structure of the ribosomal operon of *B. saltans* is similar in the closely related kinetoplastida *Leishmania*, where the ribosomal RNA genes are arranged in head- to-tail repeats (Figure 4b and Yan et al., *Molecular and biochemical parasitology* 103, no. 2 (1999): 197-210). For the cassette construction, we used only a 1 kb region of the 18S ribosomal RNA gene, which is represented in the schematic as Homologous Recombination Region 1 (HR1) and Homologous Recombination Region 2 (HR2) each, of which is 500 bp . The GFP and Neo genes are located in the middle of the construct and their expression is controlled by the presence of the *B. saltans* tubulin intergenic region (Bs TUB IGR) at the 3' and 5' ends of each gene. As the transcription in *B. saltans* is polycistronic, the presence of the tubulin intergenic region is necessary for trans-splicing and polyadenylation.

ATTACHMENTS

[SSU-GFP-Neo-SSU cassette_final.seq](#)

[18S-GFP cassette.pdf](#)

ATTACHMENTS

SSU-GFP-
Neo-SSU
cassette_fina
l.seq

18S-GFP
cassette.pdf