MAY 22, 2018

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.j9icr4e](https://doi.org/10.17504/protocols.io.j9icr4e)

External link:
<https://doi.org/10.1371/journal.pone.0196204>

Protocol Citation: Michiyo Tsuru 2018. Bone histomorphometric parameters of wild-type and knockout mice. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.j9icr4e>

MANUSCRIPT CITATION: Tsuru M, Ono A, Umeyama H, Takeuchi M, Nagata K (2018) Ubiquitin-dependent proteolysis of CXCL7 leads to posterior longitudinal ligament ossification. PLoS ONE 13(5): e0196204. doi: [10.1371/journal.pone.0196204](https://doi.org/10.1371/journal.pone.0196204)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

🌐 Bone histomorphometric parameters of wild-type and knockout mice

📁 PLOS One

Michiyo
Tsuru¹

¹Clinical Proteomics and Gene Therapy

Clinical Proteomics and Gene Therapy Laboratory

Michiyo Tsuru
Kurume University, Clinical Proteomics and Gene Therapy

ABSTRACT

The spinal bones of knockout mice were fixed in ethanol, stained by the Villanueva method, and embedded in methyl methacrylate without decalcification. The administration schedule involved a single subcutaneous injection of tetracycline administered to null and wild-type mice, followed by no administration for the next two days. Thereafter, they were given a single subcutaneous injection of calcein, followed by no administration on the next day. The results of the bone histomorphometric analysis are expressed according to the methods of the ASBMR Histomorphometry Nomenclature Committee, excluding the osteocyte parameters.

ATTACHMENTS

[プレゼンテーション1.pdf](#)

Created: Oct 11, 2017

Last Modified: May 22,
2018

PROTOCOL integer ID:
8202

1