MAR 18, 2019

🌐 bright field big patch swarming imaging

Serena Ding¹

¹Imperial College London

Behavioural Genomics

Serena Ding

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.vyhe7t6

External link:
<https://www.biorxiv.org/content/early/2018/11/01/398370>

Protocol Citation: Serena Ding 2019. bright field big patch swarming imaging. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.vyhe7t6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 28, 2018

Last Modified: Mar 18, 2019

PROTOCOL integer ID:
 18153

Keywords: behaviour, C. elegans, aggregation

ABSTRACT

For imaging swarming behaviour of 40 young adult C. elegans on a big food patch on agar using the Phoenix multi-worm tracker system. Worms are synchronised by bleaching and refeeding for 72 hours, and then 40 young adult hermaphrodites are transferred by glass pipette onto a 35 mm regular NGM plate for imaging for 20 hours at 25 fps.

BEFORE START INSTRUCTIONS

Prior to collecting the full dataset, a single batch of OP50 was grown overnight, diluted to OD600 = 0.38, aliquoted for use on each imaging day, and stored at 4°C until use.

Imaging plate preparation (Day -7)

- 1 A separate batch of imaging plates is poured exactly seven days before each imaging day and stored at 4°C.

Note

Imaging plates are 35 mm Petri dishes containing 3.5 mL low peptone (0.013% Difco Bacto) NGM agar (2% Bio/Agar, BioGene) to limit bacteria growth.

Bleach synchronising worms (Day -7 to -4)

- 2 Bleach synchronise gravid hermaphrodites. Leave on rotator at 20 °C until use.

Re-feed worms (Day -3, PM)

- 3 Re-feed starved L1 worms onto 3-4 plates using a glass pipette. Incubate at 20 °C.

Note

Culture plates are 55 mm Petri dishes containing 15 mL low peptone 0.013% Difco Bacto) NGM agar (2% Bio/Agar, BioGene), and seeded with OP50.

Seeding imaging plate (Day -1, PM)

- 4 The center of an imaging plate is seeded with a single 75 µL spot of cold diluted OP50 (OD=0.38) and allowed to inoculate overnight at room temperature before being used for imaging the next day.

Imaging (Day 0)

- 5 Wash animals off of culture plates with M9 and collect in a 15 mL Falcon tube, wash in M9 twice, and aspire as much supernatant as possible after the last wash.

- 6 Forty animals are transferred by a glass pipette onto the imaging plate in a small drop of M9, away from the bacteria lawn.
- 7 Imaging commences immediately following animal transfer in a liquid drop, on a custom-built six-camera rig (the Phoenix) equipped with Dalsa Genie cameras (G2-GM10-T2041). Twenty-hour recordings with red illumination (630 nm LED illumination, CCS Inc.) are taken at 25 Hz using Gecko software (v2.0.3.1), whilst the rig maintain the imaging plates at 20 °C throughout the recording durations.

Image data processing

- 8 Images are segmented in real time by the Gecko software.
- 9 The recordings are manually truncated post-acquisition to retain aggregation and swarming dynamics only.

Note

The start time is defined as the moment when the liquid dries and the all the worms crawl out from the initial location of the drop, and the end time is when the food is depleted and worms disperse with increased crawling speed.