JUN 29, 2021

Brucella species detection from blood and milk samples

Rania Baleela¹, Zienab Adam Ahmed Abdalh²

¹Department of Zoology, University of Khartoum; ²University of Nyala

Brucella sp detection

Rania Baleela

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bskpncvn

External link:

<http://onlinejournals.uofk.edu/index.php/SJS/article/view/1395>

Protocol Citation: Rania Baleela, Zienab Adam Ahmed Abdalh 2021. Brucella species detection from blood and milk samples.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bskpncvn>

MANUSCRIPT CITATION:

Zienab A.A. Abdallah, Rania M. H. Baleela (2017). Detection and molecular identification of Brucella species infecting Camels in Darfur, Sudan. Sudan Journal of Science, volume 9 issue 2

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Feb 20, 2021

Last Modified: Jun 29, 2021

PROTOCOL integer ID:
47471

ABSTRACT

Brucellosis is a widespread zoonotic disease which is characterized by reduced fertility and abortion in several species of animals and in human. Camels are highly susceptible to *Brucella abortu*, and *Brucella melitensis* but they are not the primary hosts.

Consumption of *Brucella* infected milk and meat from camels is a serious public health issue though. Therefore rapid detection and diagnosis of Brucellosis is crucial for any control program.

In this study, a sensitive and specific PCR assay was adapted for the detection and identification of *Brucella* spp. in 100 camels. Almost half of the collected samples were positive for *Brucella* spp. (i. e. 41%). 44% of the samples were positive for *Brucella abortus*, 43% were positive for *B. melitensis*, and 22% were positive for a co-infection of *B. abortus* and *B. melitensis*. The majority of infected camels were male adults. Most of the positive samples were collected at the slaughterhouse in Nyala. This finding is alarming as Sudanese consumers are known for eating raw camels' liver. Therefore the main recommendation of this study is to adopt this sensitive and species-specific PCR protocol for routine detection of brucellosis in animals to be slaughtered in the future .

In the mean time this protocol can be used as a confirmatory test for the detection of brucellosis in live camels as well as for other animals to be slaughtered because it is the protocol with the lowest risk for laboratory personnel.

IMAGE ATTRIBUTION

Zienab A.A. Abdallah

MATERIALS

```
<!-- /* Font Definitions */ @font-face {font-family:Calibri; panose-1:2 15 5 2 2 2 4 3 2 4; mso-font-charset:0; mso-generic-font-family:swiss; mso-font-pitch:variable; mso-font-signature:-469750017 -1073732485 9 0 511 0;} /* Style Definitions */ p.MsoNormal, li.MsoNormal, div.MsoNormal {mso-style-unhide:no; mso-style-qformat:yes; mso-style-parent:""; margin-top:0cm; margin-right:0cm; margin-bottom:10.0pt; margin-left:0cm; text-align:right; line-height:115%; mso-pagination:widow-orphan; direction:rtl; unicode-bidi:embed; font-size:11.0pt; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:"Times New Roman"; mso-fareast-theme-font:minor-fareast; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;} p.MsoListParagraph, li.MsoListParagraph, div.MsoListParagraph {mso-style-priority:34; mso-style-unhide:no; mso-style-qformat:yes; margin-top:0cm; margin-right:36.0pt; margin-bottom:10.0pt; margin-left:0cm; mso-add-space:auto; text-align:right; line-height:115%; mso-pagination:widow-orphan; direction:rtl; unicode-bidi:embed;
```

font-size:11.0pt; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:"Times New Roman"; mso-fareast-theme-font:minor-fareast; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;}

p.MsoListParagraphCxSpFirst, li.MsoListParagraphCxSpFirst, div.MsoListParagraphCxSpFirst {mso-style-priority:34; mso-style-unhide:no; mso-style-qformat:yes; mso-style-type:export-only; margin-top:0cm; margin-right:36.0pt; margin-bottom:0cm; margin-left:0cm; margin-bottom:.0001pt; mso-add-space:auto; text-align:right; line-height:115%; mso-pagination:widow-orphan; direction:rtl; unicode-bidi:embed; font-size:11.0pt; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:"Times New Roman"; mso-fareast-theme-font:minor-fareast; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;} p.MsoListParagraphCxSpMiddle, li.MsoListParagraphCxSpMiddle, div.MsoListParagraphCxSpMiddle {mso-style-priority:34; mso-style-unhide:no; mso-style-qformat:yes; mso-style-type:export-only; margin-top:0cm; margin-right:36.0pt; margin-bottom:0cm; margin-left:0cm; margin-bottom:.0001pt; mso-add-space:auto; text-align:right; line-height:115%; mso-pagination:widow-orphan; direction:rtl; unicode-bidi:embed; font-size:11.0pt; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:"Times New Roman"; mso-fareast-theme-font:minor-fareast; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;} p.MsoListParagraphCxSpLast, li.MsoListParagraphCxSpLast, div.MsoListParagraphCxSpLast {mso-style-priority:34; mso-style-unhide:no; mso-style-qformat:yes; mso-style-type:export-only; margin-top:0cm; margin-right:36.0pt; margin-bottom:10.0pt; margin-left:0cm; mso-add-space:auto; text-align:right; line-height:115%; mso-pagination:widow-orphan; direction:rtl; unicode-bidi:embed; font-size:11.0pt; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:"Times New Roman"; mso-fareast-theme-font:minor-fareast; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;} .MsoChpDefault {mso-style-type:export-only; mso-default-props:yes; font-family:"Calibri","sans-serif"; mso-ascii-font-family:Calibri; mso-ascii-theme-font:minor-latin; mso-fareast-font-family:Calibri; mso-fareast-theme-font:minor-latin; mso-hansi-font-family:Calibri; mso-hansi-theme-font:minor-latin; mso-bidi-font-family:Arial; mso-bidi-theme-font:minor-bidi;} .MsoPapDefault {mso-style-type:export-only; margin-bottom:8.0pt; line-height:107%;} @page WordSection1 {size:612.0pt 792.0pt; margin:72.0pt 72.0pt 72.0pt 72.0pt; mso-header-margin:35.4pt; mso-footer-margin:35.4pt; mso-paper-source:0;} div.WordSection1 {page:WordSection1;} /* List Definitions */ @list l0 {mso-list-id:1805459875; mso-list-type:hybrid; mso-list-template-ids:1655110872 67698703 67698713 67698715 67698703 67698713 67698715 67698703 67698713 67698715;} @list l0:level1 {mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list

l0:level2 {mso-level-number-format:alpha-lower; mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list l0:level3 {mso-level-number-format:roman-lower; mso-level-tab-stop:none; mso-level-number-position:right; text-indent:-9.0pt;} @list l0:level4 {mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list l0:level5 {mso-level-number-format:alpha-lower; mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list l0:level6 {mso-level-number-format:roman-lower; mso-level-tab-stop:none; mso-level-number-position:right; text-indent:-9.0pt;} @list l0:level7 {mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list l0:level8 {mso-level-number-format:alpha-lower; mso-level-tab-stop:none; mso-level-number-position:left; text-indent:-18.0pt;} @list l0:level9 {mso-level-number-format:roman-lower;

mso-level-tab-stop:none; mso-level-number-position:right; text-indent:-9.0pt;} ol {margin-bottom:0cm;} ul {margin-bottom:0cm;} -->

Blood, Filter paper or milk samples

NaHCO₃

NH₄Cl₆.

Distilled water

Tris-HCl

SDS

EDTA

Boric acid

Phenol

Chloroform

Isomyalcohol

Ethanol

iNtRONMaxime™ PCR premix

Primers

Agarose

Ethidium bromide

Falcon tubes

Centrifuge

pH meter

-20°C Refrigerator

Incubator.

Equipment

NanoDrop™ 3300 Fluorospectrometer

NAME

Fluorospectrometer

TYPE

NanoDrop™ 3300

BRAND

nd-3300

SKU

<https://www.thermofisher.com/order/catalog/product/ND-3300#/ND-3300> LINK

Equipment

Thermal cycler

NAME

T100 PCR thermal cycler

TYPE

BIO-RAD

BRAND

<https://www.bio-rad.com/pt-br/product/t100-thermal>

SKU

<https://www.bio-rad.com/pt-br/product/t100-thermal-cycler?ID=LGTWGIE8Z>

LINK

Expected result

specific PCR product molecular size for each Brucella species

DNA extraction: The phenol / chloroform method was used t..

- 1 Wash 1ml of blood with Red Blood Cells lysis buffer (NaHCO₃ 0.0841g, NH₄Cl 6.15135g 1000ml distilled water) twice centrifuge at 60000 rpm for 5min
- 2 Add 1ml lysis buffer and leave at 60°C overnight.
- 3 Add 500µl of Phenol/Chloroform/Isomyalcohol (25:24:1) to each sample then centrifuge for 5mn at 10000rpm ,
- 4 Transfer supernatant to another tube and add 500µl of Chloroform/Isomyalcohol (24:1)
- 5 Centrifuge at 100000rpm for 5min.
- 6 Add equal volume of 100 % ice cold ethanol and leave at -20°C overnight.
- 7 Centrifuge precipitated DNA at 14000 rpm for 10 min and wash with 70% ethanol two times.
- 8 Discard the supernatant and leave the tubes up- side-down to allow precipitated DNA to dry for 40min inside an incubator.
- 9 Dissolve the extracted DNA in 100µl double distilled water;

- 10 Quantify the concentration and store at -20°C.
- 11 Same protocol is used to extract DNA from filter paper samples: add three 1cm pieces to 500µl lysis buffer (15.76gTris-HCl (pH 8), 10% SDS, and 1.8612gEDTA in 1000ml distilled water) leave overnight at 60 °C.
- 12 Same protocol is used to extract DNA from milk samples except for the use of RBCS lysis buffer.

Detection of *Brucella* spp: For a total volume of 25 µl:

- 13 5 µl iNtRONMaxime™ PCR premix
- 14 2 µl each primer (step 15),
- 15 2 µl template DNA
- 16 *Brucella* spp. Detection primers (adopted from Khamesipouretal., 2014)

B. abortus
Forward primer: GACGAACGGAATTTTCCAATCCC
Reveres primer: TGCCGATCACTTAAGGGCCTTCAT

Expected molecular size 494bp
B. melitensis
Forward primer: AAATCGCGTCCTTGCTGGTCTGA
Reverses primer: TGCCGATCACTTAAGGGCCTTCAT
Expected molecular size 743bp

Conditions

- 17 Preheat lid to 110°C,
- 18 95°C for 5 min;
- 19 35 cycle: 95°C for min, 65 °C for 1min, 72°C for ,1 min extension at 72°C for 7 min,
- 20 Final hold at 4°C.
- 21 **Visualization:**
2% (w/v) agarose gel containing 0.5× TBE buffer (743 mM Tris–HCl (pH 8), 87 mM boric acid, and 5.3mM Na₂EDTA), stained with ethidium bromide.