JUN 19, 2019

Buffer Recipes for Protein Purification

Annie Kwon¹

¹University of California San Francisco

Annie Kwon

University of California San Francisco

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.4e8gthw

Protocol Citation: Annie Kwon 2019. Buffer Recipes for Protein Purification.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.4e8gthw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Jun 19, 2019

Last Modified: Jun 19, 2019

PROTOCOL integer ID:

24768

Keywords: Buffers, protein purification

Equilibration buffer

1

25 mM HEPES, 100 mM NaCl, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	1.461 g	2.922 g	5.844 g	11.68 8 g	23.37 6 g

Dissolve in water; bring pH up to 7.5; store at 4°C

25 mM HEPES, 500 mM NaCl, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	7.305 g	14.61 g	29.22 g	58.44 g	116.8 8 g

Dissolve in water; bring pH up to 7.5; store at 4°C

Lysis buffer

2

50 mM Na/K phosphate buffer, 500 mM NaCl, 2.5% glycerol, pH 8.0

	250 mL	500 mL	1 L	2 L	4 L
NaH ₂ PO ₄ • H ₂ O (MW=137.9923)	0.069 0 g	0.138 0 g	0.276 0 g	0.552 0 g	1.103 9 g
K ₂ HPO ₄ (MW=174.18)	2.090 2 g	4.180 3 g	8.360 7 g	16.72 13 g	33.44 26 g
NaCl (MW=58.44)	7.305 g	14.61 g	29.22 g	58.44 g	116.8 8 g
glycerol	6.25 mL	12.5 mL	25 mL	50 mL	100 mL

Dissolve in water; bring pH up to 8.0; store at 4°C

Wash buffer

3

25 mM HEPES, 1M NaCl, 10 mM imidazole, 10% glycerol, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
--	--------	--------	-----	-----	-----

HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	14.61 g	29.22 g	58.44 g	116.8 8 g	233.7 6 g
imidazole (MW=68.08)	0.170 2 g	0.340 4 g	0.680 8 g	1.361 6 g	2.723 2 g
glycerol	25 mL	50 mL	100 mL	200 mL	400 mL

Dissolve in water; bring pH up to 7.5; store at 4°C

25 mM HEPES, 1M NaCl, 10 mM imidazole, 10% glycerol, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	14.61 g	29.22 g	58.44 g	116.8 8 g	233.7 6 g
imidazole (MW=68.08)	0.340 4 g	0.680 8 g	1.361 6 g	2.723 2 g	5.446 4 g
glycerol	25 mL	50 mL	100 mL	200 mL	400 mL

Dissolve in water; bring pH up to 7.5; store at 4°C

Elution buffer

4 25 mM HEPES, 100 mM NaCl, 300 mM imidazole, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	1.461 g	2.922 g	5.844 g	11.68 8 g	23.37 6 g
imidazole (MW=68.08)	5.106 g	10.21 2 g	20.42 4 g	40.84 8 g	81.69 6 g

Dissolve in water; bring pH up to 7.5; store at 4°C

25 mM HEPES, 500 mM NaCl, 300 mM imidazole, pH 7.5

	250 mL	500 mL	1 L	2 L	4 L
HEPES (MW=238.31)	1.489 4 g	2.978 9 g	5.957 8 g	11.91 55 g	23.83 1 g
NaCl (MW=58.44)	7.305 g	14.61 g	29.22 g	58.44 g	116.8 8 g
imidazole (MW=68.08)	5.106 g	10.21 2 g	20.42 4 g	40.84 8 g	81.69 6 g

Dissolve in water; bring pH up to 7.5; store at 4°C